

American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline

Carolyn D. Runowicz, Corinne R. Leach, N. Lynn Henry, Karen S. Henry, Heather T. Mackey, Rebecca L. Cowens-Alvarado, Rachel S. Cannady, Mandi L. Pratt-Chapman, Stephen B. Edge, Linda A. Jacobs, Arti Hurria, Lawrence B. Marks, Samuel J. LaMonte, Ellen Warner, Gary H. Lyman, and Patricia A. Ganz

See accompanying article on page 539

Author affiliations appear at the end of this article.

Published online ahead of print at www.jco.org on December 7, 2015.

Support information appears at the end of this article.

Authors' disclosures of potential conflicts of interest are found in the article online at www.jco.org. Author contributions are found at the end of this article.

Additional supporting information may be found in the online version of this article.

The contents of this article are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention.

Corresponding author: Corinne R. Leach, PhD, MS, MPH, Director, Cancer and Aging Research, Behavioral Research Center, American Cancer Society, 250 Williams St, NW, Suite 600, Atlanta, GA, 30303; corinne.leach@cancer.org

© 2015 by American Cancer Society and American Society of Clinical Oncology

0732-183X/16/3406w-611w/\$20.00

DOI: 10.1200/JCO.2015.64.3809

ABSTRACT

The purpose of the American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline is to provide recommendations to assist primary care and other clinicians in the care of female adult survivors of breast cancer. A systematic review of the literature was conducted using PubMed through April 2015. A multidisciplinary expert workgroup with expertise in primary care, gynecology, surgical oncology, medical oncology, radiation oncology, and nursing was formed and tasked with drafting the Breast Cancer Survivorship Care Guideline. A total of 1,073 articles met inclusion criteria; and, after full text review, 237 were included as the evidence base. Patients should undergo regular surveillance for breast cancer recurrence, including evaluation with a cancer-related history and physical examination, and should be screened for new primary breast cancer. Data do not support performing routine laboratory tests or imaging tests in asymptomatic patients to evaluate for breast cancer recurrence. Primary care clinicians should counsel patients about the importance of maintaining a healthy lifestyle, monitor for post-treatment symptoms that can adversely affect quality of life, and monitor for adherence to endocrine therapy. Recommendations provided in this guideline are based on current evidence in the literature and expert consensus opinion. Most of the evidence is not sufficient to warrant a strong evidence-based recommendation. Recommendations on surveillance for breast cancer recurrence, screening for second primary cancers, assessment and management of physical and psychosocial long-term and late effects of breast cancer and its treatment, health promotion, and care coordination/practice implications are made.

This guideline was developed through a collaboration between the American Cancer Society and the American Society of Clinical Oncology and has been published jointly by invitation and consent in both *CA: A Cancer Journal for Clinicians* and *Journal of Clinical Oncology*. Copyright © 2015 American Cancer Society and American Society of Clinical Oncology. All rights reserved. No part of this document may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission by the American Cancer Society or the American Society of Clinical Oncology.

J Clin Oncol 34:611-635. © 2015 by American Cancer Society and American Society of Clinical Oncology

INTRODUCTION

Breast cancer is the most common noncutaneous malignancy among women, representing 4 in 10 female cancer survivors in the United States.¹ Long-term survival is common after breast cancer treatment, with a 5-year survival rate of almost 90%²; thus, addressing survivors' unique post-treatment needs is critical to providing quality health care.³

Nearly a decade ago, two landmark publications from the Institute of Medicine highlighted the importance of surveillance, health promotion, and assessing and managing the myriad of physical, psychological, spiritual, social, and practical long-term

and late effects faced by many cancer survivors after completing active treatment.^{4,5} Recent publications affirm the importance of addressing health, wellness, and quality of life (QoL) concerns of post-treatment cancer survivors.^{3,6,7} In recognition of the increasing need for information to support primary care clinicians who care for breast cancer survivors, this guideline was developed to provide recommendations to enhance the quality of clinical follow-up care for those who have completed initial treatment for female breast cancer (eg, surgery, radiation, targeted therapy, and/or chemotherapy).⁸

Although many evidence-based clinical guidelines exist for diagnosis and treatment, there are

few evidence-based clinical care guidelines addressing life-long follow-up care for survivors by cancer type. The National Comprehensive Cancer Network (NCCN) guidelines[®] are evidence and consensus based for the treatment of patients with breast cancer⁹ that include information on recommended surveillance for cancer recurrence or new cancers. The NCCN also has symptom-specific survivorship care guidelines addressing anthracycline-induced cardiac toxicity, anxiety and depression, cognitive function, fatigue, pain, sexual function, sleep disorders, healthy lifestyles and immunizations and infections.¹⁰ In addition, the American Society of Clinical Oncology (ASCO) has guidelines for the follow-up and management of patients with breast cancer^{11,12} as well as symptom-based guidelines specific to fatigue,¹³ chemotherapy-induced peripheral neuropathy (CIPN),¹⁴ and anxiety and depressive symptoms¹⁵; ASCO is also developing guidelines on the prevention and monitoring of cardiac dysfunction in survivors of adult cancers and on the management of chronic pain. Furthermore, ASCO recently endorsed the American Cancer Society (ACS) guideline on prostate cancer survivorship.¹⁶ The ACS/ASCO Breast Cancer Survivorship Care Guideline builds on prior guidelines by providing comprehensive, holistic recommendations specific to post-treatment breast cancer clinical care to help primary care clinicians better manage potential long-term and late effects and to provide timely and appropriate screening and surveillance to improve the overall health and QoL of breast cancer survivors.

This year, approximately 231,840 women will be newly diagnosed with breast cancer, and an estimated 3.1 million breast cancer survivors are alive in the United States.¹⁷ The median age at diagnosis is 61 years, and 43% are older than 65 years at diagnosis; thus, cancer survivorship must be managed in coordination with comorbidities associated with aging.¹ Approximately 61% will have localized disease, for which survival outcomes are highest (5-year relative survival rates: 99% for localized-stage breast cancer v 25% for distant-stage breast cancer).²

Breast cancer treatment depends on the stage at diagnosis, the size and location of the tumor, and tumor characteristics. Those who have stage II or III disease at diagnosis may receive more involved cancer treatment, which can result in greater likelihood and severity of the impact of treatment. Treatment generally includes two key components—treatment of the breast and local lymph nodes with surgery either with or without radiation therapy (“local therapy”) and drug treatments for cancer cells that may have spread (“adjuvant systemic therapy”) outside the breast. Surgical treatment for breast cancer includes breast-conserving surgery with radiation or mastectomy with or without radiation and with or without immediate/delayed reconstruction. In women with a very high risk of contralateral cancer from inherited susceptibility (eg, patients with mutations in the breast and ovarian cancer susceptibility genes *BRCA1/BRCA2*), contralateral prophylactic mastectomy may be performed.^{18,19} Systemic therapy may precede (“neoadjuvant”) or follow (“adjuvant”) local therapy and consists of combinations of hormonal therapy, chemotherapy, and biologic agents.

There is no standardized follow-up model for patients with early stage breast cancer who have completed surgery, chemotherapy, and radiation. Most of these women will have endocrine-responsive tumors and will require endocrine therapy for a total of 5 to 10 years. Randomized trials have shown equivalent outcomes with follow-up by either the oncologist or a primary care physician.²⁰ Shared follow-up care between one or more oncologists and the primary physician

is an additional possibility. However, the great majority of these patients will eventually be discharged back to their primary clinician for ongoing follow-up. It should be noted that these patients remain at risk indefinitely for complications of their previous cancer treatment. Most also remain at risk indefinitely for local and/or systemic recurrence of their breast cancer.

Gaps in post-treatment cancer survivorship resources and clinical follow-up care were identified through the work of the National Cancer Survivorship Resource Center (The Survivorship Center; cancer.org/survivorshipcenter), which is a collaboration between the ACS, The George Washington University (GW) Cancer Institute, and the Centers for Disease Control and Prevention (CDC) funded by a 5-year cooperative agreement from the CDC. The overarching goals of The Survivorship Center are to improve individual-level, system-level, and policy-level post-treatment survivorship clinical care, to develop resources to help survivors achieve optimal health and QoL, as well as to highlight the importance of post-treatment survivorship as a public health issue.⁸ A strategic partnership with ASCO was formed to optimize consistent, evidence-based recommendations for survivorship care in patients with breast cancer.

GUIDELINE QUESTIONS

The clinical practice guideline addresses 5 key areas of breast cancer survivorship to provide recommendations on best practice in the management of adult women after breast cancer treatment, focusing on the role of primary care clinicians and other clinicians who care for post-treatment breast cancer survivors. The 5 areas covered include 1) surveillance for breast cancer recurrence, 2) screening for second primary cancers, 3) assessment and management of physical and psychosocial long-term and late effects of breast cancer and treatment, 4) health promotion, and 5) care coordination and practice implications (Table 1).²¹⁻²³

METHODS

The methods used to develop this guideline reflect an evolving process that was influenced by ACS screening²⁴ and survivorship¹⁶ guidelines. Where appropriate, this guideline builds on the recently published ASCO symptom-based guidelines for adult cancer survivors.^{14,15,25}

PANEL FORMATION

A multidisciplinary expert workgroup was formed and tasked with drafting the Breast Cancer Survivorship Care Guideline. Workgroup members had expertise in primary care, gynecology, surgical oncology, medical oncology, radiation oncology, and nursing. In addition, a cancer survivor was included to provide a patient perspective.

LITERATURE REVIEW

The literature review began with an environmental scan of existing guidelines and guidance developed by other organizations (eg, the NCCN,^{9,10,26} ASCO,^{11,14,15,25,27} and the Children's Oncology Group *Long-Term Follow-Up Guidelines for Survivors of Childhood, Adolescent, and Young Adult Cancers* [survivorshipguidelines.org]), specific medical centers (eg, The University of Texas MD Anderson Clinical Tools and Resources Breast Cancer Survivorship algorithm,²⁸ the US Preventive Services Task Force,^{22,29,30} and the American Association of Clinical Endocrinologists³¹), individual publications³²⁻³⁵ available from other countries (eg, the Australian Cancer Survivorship Center), and publications from other expert panels (eg, the Breast Health Global Initiative guidelines³⁶).

Table 1. The Bottom Line: Recommendations for the American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline

Target population: Female adult breast cancer survivors

Target audience: Primary care providers, medical oncologists, radiation oncologists, and other clinicians caring for breast cancer survivors

Methods: An expert panel was convened to develop clinical practice guideline recommendations based on a systematic review of the medical literature

ACS/ASCO key recommendations for breast cancer survivorship care

Surveillance for breast cancer recurrence

History and physical

Recommendation 1.1: It is recommended that primary care clinicians (a) should individualize clinical follow-up care provided to breast cancer survivors based on age, specific diagnosis, and treatment protocol and as recommended by the treating oncology team (LOE = 2A); and (b) should make sure the patient receives a detailed cancer-related history and physical examination every 3 to 6 mo for the first 3 y after primary therapy, every 6-12 mo for the next 2 y, and annually thereafter (LOE = 2A).

Screening the breast for local recurrence or a new primary breast cancer

Recommendation 1.2: It is recommended that primary care clinicians (a) should refer women who have received a unilateral mastectomy for annual mammography on the intact breast and, for those with lumpectomies, an annual mammography of both breasts (LOE = 2A); and (b) should not refer for routine screening with MRI of the breast unless the patient meets high-risk criteria for increased breast cancer surveillance as per ACS guidelines (Saslow 2007²¹; LOE = 2A).

Laboratory tests and imaging

Recommendation 1.3: It is recommended that primary care clinicians should not offer routine laboratory tests or imaging, except mammography if indicated, for the detection of disease recurrence in the absence of symptoms (LOE = 2A).

Signs of recurrence

Recommendation 1.4: It is recommended that primary care clinicians should educate and counsel all women about the signs and symptoms of local or regional recurrence (LOE = 2A).

Risk evaluation and genetic counseling

Recommendation 1.5: It is recommended that primary care clinicians (a) should assess the patient's cancer family history; and (b) should offer genetic counseling if potential hereditary risk factors are suspected (eg, women with a strong family history of cancer [breast, colon, endometrial] or age 60 y or younger with triple-negative breast cancer; Moyer 2014²²; LOE = 2A).

Endocrine treatment impacts, symptom management

Recommendation 1.6: It is recommended that primary care clinicians should counsel patients to adhere to adjuvant endocrine (antiestrogen) therapy (LOE = 2A).

Screening for second primary cancers

Cancer screenings in the average-risk patient

Recommendation 2.1: It is recommended that primary care clinicians (a) should screen for other cancers as they would for patients in the general population; and (b) should provide an annual gynecologic assessment for postmenopausal women on selective estrogen receptor modulator therapies.

Assessment and management of physical and psychosocial long-term and late effects of breast cancer and treatment

Body image concerns

Recommendation 3.1: It is recommended that primary care clinicians (a) should assess for patient body image/appearance concerns (LOE = 0); (b) should offer the option of adaptive devices (eg, breast prostheses, wigs) and/or surgery when appropriate (LOE = 0); and (c) should refer for psychosocial care as indicated (LOE = IA).

Lymphedema

Recommendation 3.2: It is recommended that primary care clinicians (a) should counsel survivors on how to prevent/reduce the risk of lymphedema, including weight loss for those who are overweight or obese (LOE = 0); and (b) should refer patients with clinical symptoms or swelling suggestive of lymphedema to a therapist knowledgeable about the diagnosis and treatment of lymphedema, such as a physical therapist, occupational therapist, or lymphedema specialist (LOE = 0).

Cardiotoxicity

Recommendation 3.3: It is recommended that primary care clinicians (a) should monitor lipid levels and provide cardiovascular monitoring, as indicated (LOE = 0); and (b) should educate breast cancer survivors on healthy lifestyle modifications, potential cardiac risk factors, and when to report relevant symptoms (shortness of breath or fatigue) to their health care provider (LOE = I).

Cognitive impairment

Recommendation 3.4: It is recommended that primary care clinicians (a) should ask patients if they are experiencing cognitive difficulties (LOE = 0); (b) should assess for reversible contributing factors of cognitive impairment and optimally treat when possible (LOE = IA); and (c) should refer patients with signs of cognitive impairment for neurocognitive assessment and rehabilitation, including group cognitive training if available (LOE = IA).

Distress, depression, anxiety

Recommendation 3.5: It is recommended that primary care clinicians (a) should assess patients for distress, depression, and/or anxiety (LOE = I); (b) should conduct a more probing assessment for patients at a higher risk of depression (eg, young patients, those with a history of prior psychiatric disease, and patients with low socioeconomic status; LOE = III); and (c) should offer in-office counseling and/or pharmacotherapy and/or refer to appropriate psycho-oncology and mental health resources as clinically indicated if signs of distress, depression, or anxiety are present (LOE = I).

Fatigue

Recommendation 3.6: It is recommended that primary care clinicians (a) should assess for fatigue and treat any causative factors for fatigue, including anemia, thyroid dysfunction, and cardiac dysfunction (LOE = 0); (b) should offer treatment or referral for factors that may impact fatigue (eg, mood disorders, sleep disturbance, pain, etc) for those who do not have an otherwise identifiable cause of fatigue (LOE = I); and (c) should counsel patients to engage in regular physical activity and refer for cognitive behavioral therapy as appropriate (LOE = I).

Bone health

Recommendation 3.7: It is recommended that primary care clinicians (a) should refer post-menopausal breast cancer survivors for a baseline DEXA scan (LOE = 0); and (b) should refer for repeat DEXA scans every 2 y for women taking an aromatase inhibitor, premenopausal women taking tamoxifen and/or a GnRH agonist, and women who have chemotherapy-induced, premature menopause (LOE = 0).

(continued on following page)

Table 1. The Bottom Line: Recommendations for the American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline (continued)

Musculoskeletal health

Recommendation 3.8: It is recommended that primary care clinicians (a) should assess for musculoskeletal symptoms, including pain, by asking patients about their symptoms at each clinical encounter (LOE = 0); and (b) should offer one or more of the following interventions based on clinical indication: acupuncture, physical activity, and referral for physical therapy or rehabilitation (LOE = III).

Pain and neuropathy

Recommendation 3.9: It is recommended that primary care clinicians (a) should assess for pain and contributing factors for pain with the use of a simple pain scale and comprehensive history of the patient's complaint (LOE = 0); (b) should offer interventions, such as acetaminophen, nonsteroidal anti-inflammatory drugs, physical activity, and/or acupuncture, for pain (LOE = I); (c) should refer to an appropriate specialist, depending on the etiology of the pain once the underlying etiology has been determined (eg, lymphedema specialist, occupational therapist, etc; LOE = 0); (d) should assess for peripheral neuropathy and contributing factors for peripheral neuropathy by asking the patient about their symptoms, specifically numbness and tingling in their hands and/or feet, and the characteristics of the symptoms (LOE = 0); (e) should offer physical activity for neuropathy; and (f) should offer duloxetine for patients with neuropathic pain, numbness, and tingling (LOE = IB).

Infertility

Recommendation 3.10: It is recommended that primary care clinicians should refer survivors of childbearing age who experience infertility to a specialist in reproductive endocrinology and infertility as soon as possible (LOE = 0).

Sexual health

Recommendation 3.11: It is recommended that primary care clinicians (a) should assess for signs and symptoms of sexual dysfunction or problems with sexual intimacy (LOE = 0); (b) should assess for reversible contributing factors to sexual dysfunction and treat, when appropriate (LOE = 0); (c) should offer nonhormonal, water-based lubricants and moisturizers for vaginal dryness (LOE = IA); and (d) should refer for psychoeducational support, group therapy, sexual counseling, marital counseling, or intensive psychotherapy when appropriate (LOE = IA).

Premature menopause/hot flashes

Recommendation 3.12: It is recommended that primary care clinicians should offer selective serotonin-norepinephrine reuptake inhibitors, selective serotonin reuptake inhibitors, gabapentin, lifestyle modifications, and/or environmental modifications to help mitigate vasomotor symptoms of premature menopausal symptoms (LOE = IA).

Health promotion

Information

Recommendation 4.1: It is recommended that primary care clinicians (a) should assess the information needs of the patient related to breast cancer and its treatment, side effects, other health concerns, and available support services (LOE = 0); and (b) should provide or refer survivors to appropriate resources to meet these needs (LOE = 0).

Obesity

Recommendation 4.2: It is recommended that primary care clinicians (a) should counsel survivors to achieve and maintain a healthy weight (LOE = III); and (b) should counsel survivors if overweight or obese to limit consumption of high-calorie foods and beverages and increase physical activity to promote and maintain weight loss (LOE = IA, III).

Physical activity

Recommendation 4.3: It is recommended that primary care clinicians should counsel survivors to engage in regular physical activity consistent with the ACS guideline (Rock 2012²³) and specifically: (a) should avoid inactivity and return to normal daily activities as soon as possible after diagnosis (LOE = III); (b) should aim for at least 150 min of moderate or 75 min of vigorous aerobic exercise per wk (LOE = I, IA); and (c) should include strength training exercises at least 2 d per wk and emphasize strength training for women treated with adjuvant chemotherapy or hormone therapy (LOE = IA).

Nutrition

Recommendation 4.4: It is recommended that primary care clinicians should counsel survivors to achieve a dietary pattern that is high in vegetables, fruits, whole grains, and legumes; low in saturated fats; and limited in alcohol consumption (LOE = IA, III).

Smoking cessation

Recommendation 4.5: It is recommended that primary care clinicians should counsel survivors to avoid smoking and refer survivors who smoke to cessation counseling and resources (LOE = I).

Care coordination/practice implications

Survivorship care plan

Recommendation 5.1: It is recommended that primary care clinicians should consult with the cancer treatment team and obtain a treatment summary and survivorship care plan (LOE = 0, III).

Communication with oncology team

Recommendation 5.2: It is recommended that primary care clinicians should maintain communication with the oncology team throughout the patient's diagnosis, treatment, and post-treatment care to ensure care is evidence-based and well-coordinated (LOE = 0).

Inclusion of family

Recommendation 5.3: It is recommended that primary care clinicians should encourage the inclusion of caregivers, spouses, or partners in usual breast cancer survivorship care and support (LOE = 0).

Additional resources

More information, including a data supplement with additional evidence Tables, is available with the online version of this article at asco.org/guidelines/breast survivorship and asco.org/guidelines/wiki; patient information is available at onlinelibrary.wiley.com/doi/10.3322/caac.21319/pdf; journal-based continuing education is available at acsjournals.com/ce

Abbreviations: ACS, American Cancer Society; ASCO, American Society of Clinical Oncology; DEXA, dual-energy x-ray absorptiometry; GnRH, gonadotropin-releasing hormone; LOE, level of evidence.

LITERATURE SEARCH STRATEGY

A systematic review of the literature was conducted using PubMed through April 2015. Studies on childhood cancers, qualitative studies, and non-English publications were excluded. Search terms included: cancer survivor AND review OR meta-analysis OR systematic review OR guidelines; guidance AND breast cancer OR breast cancer survivor; breast cancer patient post-treatment AND symptom management OR late effects OR long-term

effects OR psychosocial care OR palliative care OR health promotion OR surveillance OR screening for new cancers OR self-management OR guidelines OR guidance OR follow-up OR follow-up OR adverse effects OR (chemotherapy AND adverse effects) OR (radiation AND adverse effects), OR surgery OR treatment complications OR genetic counseling and testing OR survivor or patient interventions OR provider interventions OR provider education OR barriers. Additional search attempts included breast cancer

OR breast cancer survivor OR breast cancer patient post-treatment AND (symptom-specific terms, such as lymphedema, body image, early menopause, and so on).

The highest priority was given to articles that met the following criteria: peer reviewed publication in English since 2004, unless a seminal article published before that date still carried the most weight, including randomized controlled trials (RCTs), prospective cohort studies, and well-conducted, population-based, case-control studies; large studies of more than 100 cancer cases analyzed and with high-quality assessment of covariates and analytic methods; and analyses controlled for important confounders (eg, preexisting comorbid conditions) (Table 2).

Workgroup members were also asked to consider the specific level of evidence (LOE) criteria along with consistency across studies and study designs, dose-response when presenting treatment impacts, race/ethnicity differences, and second primary cancers for which survivors are at high risk because of treatment and genetic considerations. After finalization by the workgroup, the guideline manuscript was sent to additional internal and external experts for review and comment before submission for publication. The guideline summarizes literature with the highest level of evidence (ie, RCTs). A comprehensive list of evidence is available online (see online supporting information).

This is the most recent information as of the publication date. For updates, the most recent information, and to submit new evidence, please visit asco.org/guidelines/breasturvivorship and the ASCO Guidelines Wiki (asco.org/guidelineswiki). On the basis of formal review of the emerging literature, ACS/ASCO will determine the need to update on a regular basis. At minimum, it will be updated every 5 years.

GUIDELINE DISCLAIMER

The clinical practice guidelines and other guidance published herein are provided by ACS and ASCO to assist providers in clinical decision making. The information herein should not be relied on as being complete or accurate, nor should it be considered as inclusive of all proper treatments or methods of care or as a statement of the standard of care. With the rapid development of scientific knowledge, new evidence may emerge between the time information is developed and when it is published or read. The information is not continually updated and may not reflect the most recent evidence. The information addresses only the topics specifically identified therein and is not applicable to other interventions, diseases, or stages of diseases. This information does not mandate any particular course of medical care. Furthermore, the information is not intended to substitute for the independent professional judgment of the treating provider, as the information does not account for individual variation among patients. Recommendations reflect high, moderate, or low confidence that the recommendation reflects the net effect of a given course of action. The use of words like “must,” “must not,” “should,” and “should not” indicates that a course of action is recommended or not recommended for either most or many patients, but there is latitude for the treating physician to select other

courses of action in individual cases. In all cases, the selected course of action should be considered by the treating provider in the context of treating the individual patient. Use of the information is voluntary. ACS and ASCO provide this information on an “as is” basis and make no warranty, express or suggested, regarding the information. ACS and ASCO specifically disclaim any warranties of merchantability or fitness for a particular use or purpose. ACS and ASCO assume no responsibility for any injury or damage to persons or property arising out of or related to any use of this information, or for any errors or omissions.

GUIDELINE AND CONFLICTS OF INTEREST

The expert panel was assembled in accordance with the ACS Conflict of Interest Procedures and the ASCO Conflict of Interest Management Procedures for Clinical Practice Guidelines (“Procedures”; summarized at asco.org/rwc). Members of the panel completed the ACS Guidelines Development Participant Disclosure Form, the ASCO disclosure form, and the International Committee of Medical Journal Editors Form for Disclosure of Potential Conflicts of Interest, which requires disclosure of financial and other interests that are relevant to the subject matter of the guideline, including relationships with commercial entities that are reasonably likely to experience direct regulatory or commercial impact as a result of promulgation of the guideline. Categories for disclosure include employment; leadership; stock or other ownership; honoraria, consulting or advisory role; speaker’s bureau; research funding; patents, royalties, other intellectual property; expert testimony; travel, accommodations, expenses; and other relationships. In accordance with the Procedures, the majority of the members of the panel did not disclose any such relationships.

RESULTS

In total, 1,073 articles (the list is available online; see online supporting information) met inclusion criteria; and, after full text review, 237 were included as the evidence base. Only 2% of eligible articles were rated as level I evidence, 7% were rated as level IA evidence, and 2% were rated as level IIA evidence. The majority of evidence was rated as level III (26%) and level 0 (64%). Recommendations provided in this guideline are based on current evidence in the literature and expert consensus opinion. Most of the evidence is not sufficient to warrant a strong, evidence-based recommendation. Rather, recommendations should be largely viewed as possible management strategies given the current limited evidence base and the logistical challenges of comprehensively adhering to these recommendations.

Table 2. Levels of Evidence

Level of Evidence	Criteria
I	Meta-analyses of RCTs
IA	RCT of breast cancer survivors
IB	RCT based on cancer survivors across multiple cancer sites
IC	RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing menopausal symptoms, sexual dysfunction, etc)
IIA	Non-RCTs based on breast cancer survivors
IIB	Non-RCTs based on cancer survivors across multiple sites
IIC	Non-RCTs not based on cancer survivors but on general population experiencing a specific long-term or late effect
III	Case-control study or prospective cohort study
0	Expert opinion, observational study (excluding case-control and prospective cohort studies), clinical practice, literature review, or pilot study
2A	NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Abbreviations: NCCN, National Comprehensive Cancer Network; RCTs, randomized controlled trials.

RECOMMENDATIONS

SURVEILLANCE FOR BREAST CANCER RECURRENCE

The guiding principle of surveillance is that it should consider a patient's risk of recurrence in the context of functional status and patient preferences (Table 3: Guideline for Surveillance for Breast Cancer Recurrence and Genetic Counseling). In asymptomatic patients, routine screening tests to detect recurrence are not recommended. However, a careful history is often needed to assure that patients are indeed asymptomatic. Patients at a higher risk for local recurrence should have appropriate screening.

History and Physical

Recommendation 1.1. It is recommended that primary care clinicians (a) should individualize clinical follow-up care provided to breast cancer survivors based on age, specific diagnosis, and treatment protocol as recommended by the treating oncology team (LOE = 2A); and (b) should make sure the patient receives a detailed cancer-related history and physical examination every 3 to 6 months for the first 3 years after primary therapy, every 6 to 12 months for the next 2 years, and annually thereafter by the treating oncology team (LOE = 2A).

Clinical interpretation. Perform periodic history and physical examination on all breast cancer survivors. In a previous ASCO

guideline, ASCO recommended that all female breast cancer survivors have a detailed history and physical examination every 3 to 6 months for the first 3 years after primary therapy, every 6 to 12 months for the next 2 years, and annually thereafter to detect cancer recurrence at an early stage.^{11,12} Frequency should be determined by the treating oncologist and should be based on the individual risk profile and perspective of the patient. This can be done in collaboration with the primary care clinician. Patients should be made aware of the signs and symptoms of disease recurrence and should be instructed to seek medical attention if any of the signs or symptoms occur between scheduled follow-up visits. If one or more additional oncology health care providers are following the patient, the frequency of the primary physician's visits should be adjusted accordingly. Clinicians can recommend vaccinations to their patients with a breast cancer history as appropriate based on guidelines; they can receive flu (shot, not nasal) or pneumonia vaccine at any time, including during chemotherapy, and zoster vaccine when not receiving chemotherapy.³⁷

Screening the Breast for Local Recurrence or a New Primary Breast Cancer

Recommendation 1.2. It is recommended that primary care clinicians (a) should refer women who have received a unilateral mastectomy for annual mammography on the intact breast and for those with lumpectomies an annual mammography of both breasts (LOE = 2A); and (b) *should not* refer for routine screening with

Table 3. Guideline for Surveillance for Breast Cancer Recurrence and Genetic Counseling

Recommendation	Level of Evidence ^a
It is recommended that primary care clinicians:	
Recommendation 1.1: History and physical	
(a) Should individualize clinical follow-up care provided to breast cancer survivors based on age, specific diagnosis, and treatment protocol as recommended by the treating oncology team	2A ^{b,c}
(b) Should make sure the patient receives a detailed cancer-related history and physical examination every 3-6 mo for the first 3 y after primary therapy, every 6-12 mo for the next 2 y, and annually thereafter by the treating oncology team	
Recommendation 1.2: Screening the breast for local recurrence or a new primary breast cancer	2A ^{b,c}
(a) Should refer women who have received a unilateral mastectomy for annual mammography on the intact breast and for those with lumpectomies an annual mammography of both breasts	
(b) <u>Should not</u> refer for routine screening with MRI of the breast unless the patient meets high risk criteria for increased breast cancer surveillance as per ACS guidelines	
Recommendation 1.3: Laboratory tests and imaging	2A ^{b,c}
<u>Should not</u> offer routine laboratory tests or imaging, except mammography if indicated, for the detection of disease recurrence in the absence of symptoms	
Recommendation 1.4: Signs of recurrence	2A ^{b,c}
Should educate and counsel all women about the signs and symptoms of local or regional recurrence	
Recommendation 1.5: Risk evaluation and genetic counseling	2A ^{b,c}
(a) Should assess your patient's cancer family history	
(b) Should offer genetic counseling if potential hereditary risk factors are suspected (eg, women with a strong family history of cancer [breast, colon, endometrial], or age 60 y or younger with triple-negative breast cancer)	
Recommendation 1.6: Endocrine treatment impacts, symptom management	2A ^{b,c}
Should counsel patients to adhere to adjuvant endocrine (antiestrogen) therapy	

Abbreviations: ACS, American Cancer Society; MRI, magnetic resonance imaging.

^aLevels of evidence: I indicates meta-analyses of randomized controlled trials (RCTs); IA, RCT of breast cancer survivors; IB, RCT based on cancer survivors across multiple sites; IC, RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing fatigue, lymphedema, etc); IIA, nonrandomized controlled trial (non-RCT) based on breast cancer survivors; IIB, non-RCT based on cancer survivors across multiple sites; IIC, non-RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing urinary incontinence, erectile dysfunction, etc); III, case-control or prospective cohort study; 0, expert opinion, observation, clinical practice, literature review, or pilot study. ^bNational Comprehensive Cancer Network category 2A indicates that "based upon lower-level evidence, there is uniform consensus that the intervention is appropriate." Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for NCCN Clinical Practice Guidelines in Oncology, Breast Cancer, V.2.2015. ©National Comprehensive Cancer Network, Inc 2015. All rights reserved. Accessed August 3, 2015. To view the most recent and complete version of the guideline, go online to NCCN.org. National Comprehensive Cancer Network, NCCN, NCCN Guidelines®, and all other NCCN content are trademarks owned by the National Comprehensive Cancer Network, Inc. ^cReferenced with permission from the American Society of Clinical Oncology (ASCO).

magnetic resonance imaging (MRI) of the breast unless the patient meets high risk criteria for increased breast cancer surveillance as per ACS guidelines²¹ (LOE = 2A).

Clinical interpretation. Mammography should be performed yearly on the breast treated by breast conserving surgery and on the intact contralateral breast. More frequent mammography is only warranted for evaluation or follow-up of a suspicious finding.¹¹ After mastectomy, the site of the reconstructed breast does not require imaging. While MRI of the breast is more sensitive than mammography, there is an increased risk of false-positive findings that may lead to unnecessary additional imaging and often unnecessary biopsies. The use of MRI can only be justified if the probability of missing a cancer with mammography alone is sufficiently high. In women who have not undergone bilateral mastectomy, the use of MRI of the breasts in screening for local recurrence or a new primary cancer should be restricted to women who meet the high-risk criteria of the ACS or ASCO.^{11,21,38} High risk is defined as a woman with a lifetime risk of a second primary breast cancer more than 20%, such as a woman with a *BRCA1/BRCA2* mutation or a very strong family history of breast cancer.²¹

Laboratory Tests and Imaging

Recommendation 1.3. It is recommended that primary care clinicians *should not* offer routine laboratory tests or imaging, except mammography if indicated, for the detection of disease recurrence in the absence of symptoms (LOE = 2A).

Clinical interpretation. Consistent with ASCO Guidelines and NCCN Guidelines®, routine testing with breast cancer tumor markers or imaging studies (eg, bone scan, chest x-ray, positron emission tomography-computed tomography [CT] scans, MRI scans, biomarkers) should not be performed for screening purposes, because they have not been shown to improve survival outcomes or QoL in asymptomatic patients.^{9,11} Chest x-rays and advanced body imaging (eg, CT, MRI, positron emission tomography-CT, bone scan) should be ordered only if disease recurrence is suspected.³⁹ Randomized trials in the 1980s that compared clinical follow-up with periodic advanced imaging did not demonstrate a survival advantage with advanced imaging and did show a significant rate of false-positive findings with additional testing.⁴⁰⁻⁴²

Signs of Recurrence

Recommendation 1.4. It is recommended that primary care clinicians should educate and counsel all women about the signs and symptoms of local or regional recurrence (LOE = 2A).

Clinical interpretation. Physicians should educate and counsel patients about the signs and symptoms of local or regional recurrence, including new lumps (eg, in underarm or neck), rash or skin changes on the breast or chest wall, chest pain, changes in the contour/shape/size of the breast, and swelling of the breast or arm.¹¹ Evaluation of patient-reported symptoms is essential in detecting a recurrence as early as possible, which may impact survival.

Risk Evaluation and Genetic Counseling

Recommendation 1.5. It is recommended that primary care clinicians (a) should assess the patient's cancer family history; and (b) should offer genetic counseling if potential hereditary risk factors are suspected (eg, women with a strong family history of cancer

[breast, colon, endometrial], or age 60 years or younger with triple-negative breast cancer)²² (LOE = 2A).

Clinical interpretation. To identify those women with breast cancer who have a high risk of a second primary breast cancer and/or may have a genetic susceptibility to cancer that may affect other family members, a detailed history, including key risk factors and paternal and maternal family history, should be obtained for all patients. Those with a family history of breast or ovarian cancer or with cancer in a certain age group and/or cancer type should be referred for genetic counseling for consideration of testing for hereditary predisposition to genetic mutations. Specifically, genetic counseling for consideration of testing for hereditary predisposition to gene mutations should be recommended for breast cancer survivors with in the following characteristics: those with at least one grandparent of Ashkenazi Jewish heritage, younger than age 50 years at diagnosis, with a history of ovarian cancer at any age or in any first-degree or second-degree relative, with a first-degree relative who had breast cancer diagnosed before age 50 years, with two or more first-degree or second-degree relatives diagnosed with breast cancer at any age, with a diagnosis of bilateral breast cancer, with a history of breast cancer in a male relative, or any survivor diagnosed at age 60 years or younger with triple-negative breast cancer.²² It is important to periodically review these issues with the patient, because some survivors may not have been offered genetic counseling or testing at the time of diagnosis, and/or new cancer events may have occurred in the family after the initial diagnosis and treatment. Because new primary cancers may be associated with some hereditary syndromes, identifying the risk of genetic mutations in the survivor may help to formulate a prevention strategy to reduce the risk of a new cancer. In addition, this information also could be helpful to family members.

Genetic testing should be preceded by consultation with a genetics counselor or other trained professional to assure full discussion of the risks and benefits and to assure that other genetic syndromes beyond *BRCA1* and *BRCA2* breast/ovarian syndromes are considered. Depending on the hereditary gene that is identified, different screening and prevention strategies may be offered. Recommendations for alternate screening and prevention strategies depend on the specific genetic syndrome and should be left to a trained professional in coordination with the oncology team and the primary care clinician.

Endocrine Treatment Impacts, Symptom Management

Recommendation 1.6. It is recommended that primary care clinicians should counsel patients to adhere to adjuvant endocrine (antiestrogen) therapy (LOE = 2A).

Clinical interpretation. Endocrine therapy (tamoxifen, aromatase inhibitors, or ovarian suppression therapy) used as adjuvant systemic therapy for 5 to 10 years reduces the risk of recurrence and of subsequent second primary breast cancers and improves overall survival. Adherence to endocrine therapy is necessary to achieve its survival benefits. Unfortunately, some women discontinue endocrine therapy because of cost, adverse effects, and other reasons. Reported adherence to a 5-year course of therapy ranges from 50% to 92% of breast cancer patients.⁴³ Primary care clinicians should assess and encourage adherence to adjuvant endocrine therapy at each visit.⁹

SCREENING FOR SECOND PRIMARY CANCERS

Cancer Screenings in the Average-Risk Patient

Recommendation 2.1. It is recommended that primary care clinicians (a) should screen for other cancers, as they would for patients in the general population; and (b) should provide an annual gynecologic assessment for postmenopausal women on selective estrogen receptor modulator therapies (SERMs) (Table 4).

Clinical interpretation. Women should be advised to follow the ACS screening and early detection guidelines for cervical, colorectal, endometrial, and lung cancers detailed in Table 4.⁴⁴ Postmenopausal women who are taking SERMs, such as tamoxifen, should be advised to report any vaginal spotting or bleeding, because these drugs slightly increase the risk of endometrial cancer in postmenopausal women. In the absence of abnormal vaginal spotting or bleeding, periodic imaging is not of value and may lead

to unwarranted biopsies.⁴⁵ Discuss the risks, benefits and limitations of screening modalities with your patients.

ASSESSMENT AND MANAGEMENT OF PHYSICAL AND PSYCHOSOCIAL LONG-TERM AND LATE EFFECTS OF BREAST CANCER AND TREATMENT

The risk of physical long-term and late effects after therapy for breast cancer is associated with several factors, including: (a) type of treatment, (b) duration and dose of treatment(s) (increasing cumulative dose and duration of therapy increase the potential risk), (c) specific type of chemotherapy, (d) receipt of and type of hormone treatment, and (e) age of patient during treatment. Modalities of treatment include surgery, radiation therapy, chemotherapy, targeted therapy, and/or endocrine therapy. Primary care clinicians should refer to the patient's cancer treatment summary, if available, for specific drugs and doses (see Recommendation 5.1). Table 5 lists

Table 4. American Cancer Society Guidelines for the Early Detection of Cancer in Average-Risk, Asymptomatic Individuals^a

Cancer Site	Population	Test or Procedure	Frequency
Cervix	Women, ages 21-65 y	Pap test and HPV DNA test	Cervical cancer screening should begin at age 21 y; for women ages 21-29 y, screening should be done every 3 y with conventional or liquid-based Pap tests; for women ages 30-65 y, screening should be done every 5 y with both the HPV test and the Pap test (preferred) or every 3 y with the Pap test alone (acceptable); women older than 65 y who have had ≥ 3 consecutive negative Pap tests or ≥ 2 consecutive negative HPV and Pap tests within the last 10 y, with the most recent test occurring within the last 5 y, and women who have had a total hysterectomy should stop cervical cancer screening if they no longer have a cervix and are without a history of cervical intraepithelial lesion grade 2 or a more severe diagnosis in the past 20 y or if they have ever had cervical cancer; women at any age should not be screened annually by any screening method
Colorectal	Women aged 50 y and older	FOBT with at least 50% test sensitivity for cancer, or FIT with at least 50% test sensitivity for cancer, or Stool DNA test, ^b or FSIG, or DCBE, or Colonoscopy CT colonography	Annual, starting at age 50 y; testing at home with adherence to manufacturer's recommendation for collection techniques and number of samples is recommended; FOBT with the single stool sample collected on the clinician's fingertip during a DRE in the health care setting is not recommended; guaiac-based toilet bowl FOBT tests also are not recommended; compared with guaiac-based tests for the detection of occult blood, immunochemical tests are more patient-friendly and are likely to be equal or better in sensitivity and specificity; there is no justification for repeating FOBT in response to an initial positive finding Interval uncertain, starting at age 50 y Every 5 y, starting at age 50 y; FSIG can be performed alone or consideration can be given to combining FSIG performed every 5 y with a highly sensitive guaiac-based FOBT or FIT performed annually Every 5 y, starting at age 50 y Every 10 y, starting at age 50 y Every 5 y, starting at age 50 y
Endometrial	Women, at menopause		At the time of menopause, women at average risk should be informed about the risks and symptoms of endometrial cancer and strongly encouraged to report any unexpected bleeding or spotting to their physicians
Lung	Current or former smokers ages 55-74 y in good health with at least a 30 pack-y history	LDCT	Clinicians with access to high-volume, high-quality lung cancer screening and treatment centers should initiate a discussion about lung cancer screening with apparently healthy patients ages 55-74 y who have at least a 30 pack-y smoking history and who currently smoke or have quit within the past 15 y; a process of informed and shared decision-making with a clinician related to the potential benefits, limitations, and harms associated with screening for lung cancer with LDCT should occur before any decision is made to initiate lung cancer screening; smoking-cessation counseling remains a high priority for clinical attention in discussions with current smokers, who should be informed of their continuing risk of lung cancer; screening should not be viewed as an alternative to smoking cessation
Cancer-related checkup	Women, aged 20 y and older		On the occasion of a periodic health examination, the cancer-related checkup should include examination for cancers of the thyroid, ovaries, lymph nodes, oral cavity, and skin as well as health counseling about tobacco, sun exposure, diet and nutrition, risk factors, sexual practices, and environmental and occupational exposures

Abbreviations: CT, computed tomography; DCBE, double-contrast barium enema; DRE, digital rectal examination; FIT, fecal immunochemical test; FOBT, fecal occult blood test; FSIG, flexible sigmoidoscopy; HPV, human papillomavirus; LDCT, low-dose helical computed tomography; Pap, Papanicolaou.

^aRecommendation 2.1: Among average-risk patients, it is recommended that primary care clinicians (a) should screen for other cancers as they would for patients in the general population; and (b) should provide an annual gynecological assessment for postmenopausal women on selective estrogen receptor modulator therapies. ^bThe stool DNA test approved for colorectal cancer screening in 2008 is no longer commercially available; new stool DNA tests are presently undergoing evaluation and may become available at some future time.

Table 5. Summary of Long-Term and Late Effects by Treatment Type^a

Treatment Type	Long-Term Effects	Late Effects
Surgery	<ul style="list-style-type: none"> • Lack of skin sensitivity • Body image issues • Sexual dysfunction • Numbness • Pain • Limited range of motion • Weakness • Poor cosmetic outcome 	<ul style="list-style-type: none"> • Lymphedema • Neuropathy
Radiation therapy to the breast/ chest wall/regional lymph nodes	<ul style="list-style-type: none"> • Fatigue^{b,c} • Skin sensitivity/pain • Sexual dysfunction • Pain • Pneumonitis^{b,c} • Poor cosmetic outcome • Breast atrophy/asymmetrical breast volume • Lymphedema^b • Numbness or weakness of the upper extremity^b 	<ul style="list-style-type: none"> • Skin discoloration • Breast may be slightly smaller and firmer than the nonirradiated side (breast asymmetry) • Skin sensitivity/pain • Telangiectasia • Sexual dysfunction • Lymphedema^b • Shortness of breath (lung pneumonitis or fibrosis)^{b,c} • Cardiovascular disease (eg, pericardial effusion, pericarditis)^c • Numbness or weakness of the upper extremity^{c,d} • Second primary cancers (eg, soft-tissue sarcomas of thorax, shoulder, and pelvis; lung cancer)^{b,c}
Chemotherapy	<ul style="list-style-type: none"> • Cognitive impairment • Fatigue • Ovarian failure with or without menopausal symptoms • Sexual dysfunction • Change in libido • Infertility • Weight gain • Obesity • Neuropathy, especially after taxanes • Oral health issues • Hair loss 	<ul style="list-style-type: none"> • Osteoporosis/osteopenia • Increased risk of cardiovascular disease (cardiomyopathy, congestive heart failure) with anthracycline-based chemotherapy • Increased risk of leukemia and myelodysplastic syndrome with alkylating agents, anthracyclines, other topoisomerase II inhibitors, and other agents with immunosuppressive potential
Hormonal therapy Tamoxifen	<ul style="list-style-type: none"> • Hot flushes • Changes in menstruation • Mood changes • Increased triglycerides • Vaginal dryness • Decreased libido • Musculoskeletal symptoms/pain • Cholesterol elevation 	<ul style="list-style-type: none"> • Increased risk of stroke • Increased risk of endometrial cancer • Increased risk of blood clots • Osteopenia in premenopausal women • Increased risk of osteoporosis • Increased risk of fractures
Aromatase inhibitors	<ul style="list-style-type: none"> • Musculoskeletal symptoms/pain • Cholesterol elevation 	
Targeted therapy Trastuzumab	<ul style="list-style-type: none"> • Increased risk of cardiac dysfunction 	
General psychosocial long-term and late effects	<ul style="list-style-type: none"> • Depression • Distress—multifactorial unpleasant experience of psychological, social, and/or spiritual nature • Worry, anxiety • Fear of recurrence • Fear of pain • End-of-life concerns: Death and dying • Loss of sexual function and/or desire • Challenges with body image • Challenges with self-image • Relationship and other social role difficulties • Return-to-work concerns and financial challenges 	

^aNote: Long-term effect is something that starts during treatment and does not subside, such as pain, fatigue, cognitive changes. In contrast, a late effect is something that develops much later, such as a second malignancy, heart failure, lymphedema. Some long-term effects are the same or overlap with late effects; for example, lymphedema can be either. ^bRisks are increased in patients who also received radiotherapy to the supraclavicular lymph nodes. ^cRisks are increased in patients who also received radiotherapy to the internal mammary lymph nodes. ^dThere is a need to be careful, because these can also be signs of recurrent cancer, typically with pain; an appropriate consultation with the radiation oncologist may be warranted.

potential physical and psychosocial long-term and late effects associated with surgery, radiation, chemotherapy, hormone therapy, and targeted treatment. Long-term effects are medical problems that develop during active treatment and persist after the completion of treatment, whereas late effects are medical problems that develop or become apparent months or years after treatment is completed. Recommendations for the assessment and management of specific physical and psychosocial long-term and late effects most commonly experienced by breast cancer survivors are detailed in [Table 6](#).

Body Image Concerns

Recommendation 3.1. It is recommended that primary care clinicians (a) should assess for patient body image/appearance concerns (LOE = 0); (b) should offer the option of adaptive devices (eg, breast prostheses, wigs) and/or surgery when appropriate (LOE = 0); and (c) should refer for psychosocial care as indicated (LOE = IA).

Clinical interpretation. Body image/appearance changes can be a major area of concern, affecting from 31% to 67% of breast cancer survivors.⁴⁶ Factors such as the loss of a breast, scarring and/or lymphedema after surgery, hair loss or sexual dysfunction/chemotherapy-related early menopause, skin changes from radiation, and weight gain can all lead to changes in body image with negative implications on short-term and long-term QoL.⁴⁷ These body image implications are especially relevant for young breast cancer survivors.⁴⁷⁻⁵⁰ Among female breast cancer survivors who are sexually active, greater body image problems have been associated with mastectomy with or without reconstruction, hair loss from chemotherapy, concern with weight change, decreased self-esteem before the time of cancer diagnosis, poorer mental health, and a partner's difficulty understanding one's feelings.⁴⁹

Patients with radiation-associated breast/soft-tissue fibrosis should be considered for therapy with oral pentoxifylline (Trental) and vitamin E.⁵¹⁻⁵⁴ A radiation oncologist should evaluate such patients to assure that the clinical findings are consistent with the radiation treatment delivered. Recurrent cancer or infection can sometimes be mistaken for radiation-associated fibrosis. Primary care clinicians should refer back to the oncologist (medical, surgical, or radiation) if they ever are uncertain about potential recurrence.

Breast-reconstructive surgery should be considered in women who do not feel comfortable with the results of their initial breast surgery, whether lumpectomy or mastectomy, because there may be ways to improve symmetry or appearance. Breast forms/prostheses or bras are options that are available to survivors who have undergone mastectomy or non-breast-conserving therapies and generally require a prescription from a physician. Special lingerie, attachable nipples, and other resources can help the survivor regain self-esteem. Most insurance plans cover mastectomy prostheses and specialty bras. In addition, wigs, hats, scarves, and other accessories are available for survivors who experience permanent hair loss (alopecia is a rare complication of chemotherapy) or changes in hair color or texture. Women should check with their insurance company and obtain a prescription from their physicians. Breast cancer care centers or specialists often maintain directories of resources in their community where patients can find these products.

Primary care clinicians should refer patients who have body image concerns that are not corrected by the above options for psychosocial care. For example, for women who experience early

menopause or for those with sexual concerns, support groups, psychotherapy, cognitive behavioral therapy, couple-based interventions, or sex therapy may be helpful. Couple-based interventions can promote coping skills and provide specific techniques to address body image issues.⁵⁵

Lymphedema

Recommendation 3.2. It is recommended that primary care clinicians (a) should counsel survivors on how to prevent/reduce risk of lymphedema, including weight loss for those who are overweight or obese (LOE = 0); and (b) should refer patients with clinical symptoms or swelling suggestive of lymphedema to a therapist knowledgeable about the diagnosis and treatment of lymphedema, such as a physical therapist, occupational therapist, or lymphedema specialist (LOE = 0).

Clinical interpretation. All breast cancer survivors who undergo breast surgery and/or radiation are at risk for lymphedema. This is arm, breast, or chest wall swelling as a result of a blockage of the lymphatic fluid from the arm and/or breast, leading to retention of fluid and swelling. The incidence of lymphedema among breast cancer survivors varies widely,⁵⁶ although it is estimated that over 40% of survivors will experience lymphedema to some degree.^{57,58} The risk of lymphedema is much lower with sentinel lymph node dissections than with the full axillary lymph node dissection previously performed in all cases.⁵⁹ Lymphedema may occur immediately after treatment or develop after many years. Radiation treatment may cause or exacerbate lymphedema, especially radiation to the supraclavicular lymph nodes or axilla.⁶⁰

The degree of swelling associated with lymphedema varies widely, even for those who receive similar surgery and/or radiation. In most cases, it is generally limited in extent and is not disabling. In some cases, the swelling is extensive and leads to significant disability, such as limitation of the ability to perform fine motor functions with the hand as well as limited range of motion of other affected joints.⁶¹ In addition, the swelling may cause symptoms ranging from mild discomfort to overt pain. The swelling may be of a noticeable degree, making clothes difficult to fit and causing pain from the added weight of the arm. The International Society of Lymphology provides a staging system to categorize the extent of lymphedema.⁶² Patients with lymphedema are also at greater risk for the development of cellulitis of the breast, arm, or chest, which, especially if not promptly treated with antibiotics, may exacerbate lymphedema.

More studies are needed in the area of lymphedema prevention above and beyond biopsy types and surgical strategies (sentinel lymph node dissections and axillary reverse mapping).⁶³ Obesity is also a risk factor for lymphedema,^{64,65} so physicians should recommend weight loss for those who are overweight or obese and emphasize the importance of maintaining a normal weight (see also Recommendation 5.2). Historically, patients with axillary lymphadenectomies/radiation have been advised to avoid physical activity and heavy lifting with the arm on the affected side. However, one study has shown that supervised, slowly progressive resistance training is safe and effective for breast cancer survivors with regard to lymphedema development. Furthermore, this type of physical activity may reduce the likelihood of arm swelling among breast cancer survivors at high risk for lymphedema who have had five or more lymph nodes removed and may improve the symptoms and severity for those in whom the condition was already present.⁶⁶ While

Table 6. Guideline for Assessment and Management of Physical and Psychosocial Long-Term/Late Effects

Recommendation	Level of Evidence ^a
It is recommended that primary care clinicians:	
Recommendation 3.1: Body image concerns	
(a) Should assess for patient body image/appearance concerns	0 (assessment)
(b) Should offer the option of adaptive devices (eg, breast prostheses, wigs) and/or surgery when appropriate	0 (adaptive devices)
(c) Should refer for psychosocial care as indicated	IA (couple-based intervention)
Recommendation 3.2: Lymphedema	
(a) Should counsel survivors on how to prevent/reduce the risk of lymphedema, including weight loss for those who are overweight or obese	0 (prevention)
(b) Should refer patients with clinical symptoms or swelling suggestive of lymphedema to a therapist knowledgeable about the diagnosis and treatment of lymphedema, such as a physical therapist, occupational therapist, or lymphedema specialist	0 (referral)
Recommendation 3.3: Cardiotoxicity	
(a) Should monitor lipid levels and provide cardiovascular monitoring, as indicated	0 (monitoring)
(b) Should educate breast cancer survivors on healthy lifestyle modifications, potential cardiac risk factors, and when to report relevant symptoms (shortness of breath or fatigue) to their health care provider	I (lifestyle modifications)
Recommendation 3.4: Cognitive impairment	
(a) Should ask patients if they are experiencing cognitive difficulties	0 (assessment)
(b) Should assess for reversible contributing factors of cognitive impairment and optimally treat when possible	IA (contributing factors)
(c) Should refer patient with signs of cognitive impairment for neurocognitive assessment and rehabilitation, including group cognitive training if available	IA (group cognitive rehabilitation)
Recommendation 3.5: Distress, depression, and anxiety	
(a) Should assess patients for distress, depression, and/or anxiety	I (assessment)
(b) Should conduct a more probing assessment for patients at a higher risk of depression (eg, young patients, those with a history of prior psychiatric disease, and patients with low socioeconomic status)	III (at-risk groups)
(c) Should offer in-office counseling and/or pharmacotherapy and/or refer to appropriate psycho-oncology and mental health resources as clinically indicated if signs of distress, depression, or anxiety are present	I (interventions)
Recommendation 3.6: Fatigue	
(a) Should assess for fatigue and treat any causative factors for fatigue, including anemia, thyroid dysfunction, and cardiac dysfunction	0 (assessment)
(b) Should offer treatment or referral for factors that may impact fatigue (eg, mood disorders, sleep disturbance, pain, etc) for those who do not have an otherwise identifiable cause of fatigue	I (causative factors)
(c) Should counsel patients to engage in regular physical activity and refer for cognitive behavioral therapy (CBT) as appropriate	I (physical activity, CBT)
Recommendation 3.7: Bone health	
(a) Should refer postmenopausal breast cancer survivors for a baseline DEXA scan	0 (baseline DEXA)
(b) Should refer for repeat DEXA scans every 2 y for women taking an aromatase inhibitor, premenopausal women taking tamoxifen and/or a GnRH agonist, and women who have chemotherapy-induced premature menopause	0 (high risk)
Recommendation 3.8: Musculoskeletal health	
(a) Should assess for musculoskeletal symptoms, including pain, by asking patients about their symptoms at each clinical encounter	0 (assessment)
(b) Should offer one or more of the following interventions based on clinical indication: acupuncture, physical activity, and referral for physical therapy or rehabilitation	III (interventions)
Recommendation 3.9: Pain and neuropathy	
(a) Should assess for pain and contributing factors for pain with the use of a simple pain scale and comprehensive history of the patient's complaint	0 (assessment)
(b) Should offer interventions, such as acetaminophen, nonsteroidal anti-inflammatory drugs, physical activity, and/or acupuncture, for pain	I (interventions)
(c) Should refer to an appropriate specialist, depending on the etiology of the pain once the underlying etiology has been determined (eg, lymphedema specialist, occupational therapist, etc)	0 (referral)
(d) Should assess for peripheral neuropathy and contributing factors for peripheral neuropathy by asking the patient about their symptoms, specifically numbness and tingling in their hands and/or feet, and the characteristics of those symptoms	0 (assessment)
(e) Should offer physical activity for neuropathy.	IA (physical activity)
(f) Should offer duloxetine for patients with neuropathic pain, numbness, and tingling	IB (duloxetine)
Recommendation 3.10: Infertility	
Should refer survivors of childbearing age who experience infertility to a specialist in reproductive endocrinology and infertility as soon as possible	0
Recommendation 3.11: Sexual health	
(a) Should assess for signs and symptoms of sexual dysfunction or problems with sexual intimacy	0 (assessment)
(b) Should assess for reversible contributing factors to sexual dysfunction and treat when appropriate	0 (contributing factors)

(continued on following page)

Table 6. Guideline for Assessment and Management of Physical and Psychosocial Long-Term/Late Effects (continued)

Recommendation	Level of Evidence ^a
(c) Should offer nonhormonal, water-based lubricants and moisturizers for vaginal dryness	IA (nonhormonal lubricants and moisturizers)
(d) Should refer for psychoeducational support, group therapy, sexual counseling, marital counseling or intensive psychotherapy, when appropriate	IA (counseling)
Recommendation 3.12: Premature menopause/hot flashes Should offer selective serotonin-norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), gabapentin, lifestyle modifications and/or environmental modifications to help mitigate vasomotor symptoms of premature menopausal symptoms	IA (SNRI and SSRI use)

Abbreviations: DEXA indicates dual-energy x-ray absorptiometry; GnRH, gonadotropin-releasing hormone.
^aLevel of evidence: I indicates meta-analyses of randomized controlled trials (RCTs); IA, RCT of breast cancer survivors; IB, RCT based on cancer survivors across multiple sites; IC, RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing fatigue, lymphedema, etc); IIA, nonrandomized controlled trial (non-RCT) based on breast cancer survivors; IIB, non-RCT based on cancer survivors across multiple sites; IIC, non-RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing urinary incontinence, erectile dysfunction, etc); III, case-control or prospective cohort study; O, expert opinion, observation, clinical practice, literature review, or pilot study.

the results of this study suggest a promising intervention for lymphedema, additional research is warranted. Therefore, primary care clinicians should focus on the early identification and management of lymphedema among their patients with breast cancer.⁶⁷

Patients who develop clinical symptoms or swelling suggestive of lymphedema should be referred to a therapist knowledgeable about the diagnosis and treatment of lymphedema. Depending on available resources in a community, this may be a physical⁶⁸ or occupational⁶⁹ therapist or a specialist therapist trained in lymphedema management.

Cardiotoxicity

Recommendation 3.3. It is recommended that primary care clinicians (a) should monitor lipid levels and provide cardiovascular monitoring, as indicated (LOE = 0); and (b) should educate breast cancer survivors on healthy lifestyle modifications, potential cardiac risk factors, and when to report relevant symptoms (shortness of breath or fatigue) to their health care provider (LOE = I).

Clinical interpretation. Radiation, chemotherapy, and hormonal/endocrine therapy with aromatase inhibitors have been associated with an increased risk of cardiovascular disease in patients with breast cancer.^{70,71} The risk of heart disease increases in postmenopausal women, as endogenous estrogens in younger women contribute to the low prevalence of cardiovascular disease in that population. Therefore, breast cancer patients who experience treatment-related early menopause may be at higher risk for heart disease than age-matched women in the general population. The chemotherapeutic agents epirubicin and doxorubicin are associated with a low but real risk of cardiomyopathy⁷²; similarly, trastuzumab is associated with an increased risk of cardiac dysfunction, most notably when given concomitantly or after an anthracycline, as has been detailed by the European Society for Medical Oncology.⁷³ Aromatase inhibitors can raise cholesterol levels^{74,75} and the risk of diabetes. Significant weight gain may lead to hypertension and insulin resistance, which further elevate the risk of cardiovascular disease. Primary care clinicians should monitor lipid levels and engage in cardiovascular monitoring consistent with clinical standards for other high-risk populations (eg, US Preventive Services Task Force recommendations²⁹).

It is important to educate breast cancer survivors about lifestyle modifications, including smoking cessation, diet, and exercise (see Recommendations 4.3-4.5), that may reduce the risk or severity of

cardiotoxicity or cardiovascular diseases in general. Patients should be advised to be aware of the potential risk for cardiotoxicities and report symptoms such as shortness of breath or fatigue (without an otherwise identifiable cause) to their health care provider. However, routine screening or testing for cardiovascular disease in asymptomatic patients beyond careful history and physical examination are not warranted. ASCO is currently developing a guideline on the prevention and monitoring of cardiac dysfunction in survivors of adult cancers.

Cognitive Impairment

Recommendation 3.4. It is recommended that primary care clinicians (a) should ask patients if they are experiencing cognitive difficulties (LOE = 0); (b) should assess for reversible contributing factors of cognitive impairment and optimally treat when possible (LOE = IA); and (c) should refer patients with signs of cognitive impairment for neurocognitive assessment and rehabilitation, including group cognitive training if available (LOE = IA).

Clinical interpretation. Impairment in cognitive function as a result of cancer and its treatment can lead to distress and impaired QoL in breast cancer survivors. Up to 75% of breast cancer patients in treatment and 35% after treatment report cognitive impairment, including problems with concentration, executive function, and memory.⁷⁶⁻⁷⁸ Cognitive impairment can also have detrimental effects on the survivor's role within the family, in the workplace, and in society.^{79,80} Clinicians should ask patients if they are having cognitive difficulties and listen to family members' reporting of patient cognitive symptoms. If this is an issue, consultation with a neuropsychologist for assessment and referral for cognitive rehabilitation strategies should be offered as one would do with the general population.

The causes of cognitive impairment are thought to be multifactorial and may include treatable conditions, such as fatigue, insomnia, and depression. Links have been suggested between cognitive impairment and adjuvant chemotherapy,⁸¹⁻⁸⁴ surgery/anesthesia, endocrine therapy,^{76,85} as well as cancer itself.⁷⁶ Research suggests that older adults and/or those with lower cognitive reserves may be more susceptible to cognitive impairment from treatment⁸⁶ and that there may also be genetic factors associated with certain estrogen metabolism genes.^{87,88}

The treatment of cognitive impairment in the breast cancer survivor is not well established. A few studies have shown some success with pharmaceuticals, such as modafinil,⁸⁹ that have helped

improve cognitive impairment, but the data are not consistent.⁹⁰ Cognitive rehabilitation strategies, including the practice of group cognitive training (ie, interventions geared toward improving, restoring, or maintaining mental function through structured, repetitive practice of tasks posing a mental challenge or requiring the person to problem solve) has been found to be helpful at reducing cognitive impairment in breast cancer survivors.^{91,92} Primary care clinicians should provide referral for neurocognitive assessment and rehabilitation as clinically indicated.

Distress, Depression, and Anxiety

Recommendation 3.5. It is recommended that primary care clinicians (a) should assess patients for distress, depression, and/or anxiety (LOE = I); (b) should conduct a more probing assessment for patients at a higher risk of depression (eg, young patients, those with a history of prior psychiatric disease, and patients with low socioeconomic status; LOE = III); and (c) should offer in-office counseling and/or pharmacotherapy and/or refer to appropriate psycho-oncology and mental health resources as clinically indicated if signs of distress, depression, or anxiety are present (LOE = I).

Clinical interpretation. Many cancer survivors report ongoing difficulties in recovery and returning to “normal” after treatment.^{1,2,17,93} Some survivors of cancer experience fear of recurrence (FOR),⁹⁴ which contributes to significant mental health problems for which they may already have an increased risk, including distress, depression, and anxiety.^{95,96} African American breast cancer survivors and those who are older have been found to have less FOR. A shorter interval of time since diagnosis, having received chemotherapy, and having more symptoms, especially pain, have been related to higher levels of FOR.⁹⁷ Prevalence estimates for anxiety, depression, and distress in cancer survivors vary widely as a result of inconsistency in the use of measurement tools and differences in methodological approaches, such as the choice of comparators from the general population. The estimated prevalence of anxiety and depression is 17.9% and 11.6%, respectively, among the general cancer survivor population.⁹⁸ In a systematic review of observational studies, the prevalence of depression and anxiety specifically among breast cancer survivors was 22% (range, 13% to 56%) using the Center for Epidemiologic Studies for Depression, 22% (range, 17% to 48%) using the Beck Depression Inventory, and 10% (range, 1% to 22%) using the Hospital Anxiety and Depression Scale.⁹⁹

To provide timely and appropriate support for their patients with a history of breast cancer, primary care clinicians should be familiar with the mental health concerns they may experience, the tools to screen for and assess these problems, and the resources to care for their patients. A tool for initial screening is the distress thermometer (NCCN.org), with scores from 0 (no distress) to 10 (extreme distress). A score of 4 or higher¹⁰⁰ suggests a level of distress that has clinical significance. The Patient Health Questionnaire-9¹⁰¹ and the Generalized Anxiety Disorder 7-item scale¹⁰² are validated methods for screening for depression and anxiety, respectively, and are available free online at phqscreener.com/. For patients with elevated scores on these screeners, further discussion and assessment of the issues are needed, as in the general population. For more information on screening and assessing adults with cancer for psychosocial distress, depression and anxiety algorithms reproduced from the ASCO guidelines are provided online (see online supporting information).¹⁵

The risk of having major depression after a diagnosis of breast cancer was higher among younger patients, patients with a history of prior psychiatric disease, patients with low socioeconomic status, and those who were unemployed.¹⁰³ Among patients with breast cancer, decreased libido, poor self-image, and relationship issues were common among those who were depressed.¹⁰⁴⁻¹⁰⁶

Treatment for depression and anxiety in cancer patients and survivors with medication and psychotherapy is comprehensively described in the recent ASCO anxiety and depression guideline adaptation (instituteofquality.org/screening-assessment-and-care-anxiety-and-depressive-symptoms-adults-cancer-american-society).¹⁵ In addition to or instead of pharmacotherapy, mindfulness-based approaches, expression of positive emotions, spiritual interventions, hope therapy, and meaning-making interventions have shown promise in addressing psychosocial needs of breast cancer survivors.¹⁰⁷ Although the methodology used to study the effectiveness of these interventions varied, survivors experienced positive changes, such as enhanced QoL and well-being.¹⁰⁷ If a patient has a clinically significant score on any of the previously discussed instruments, it is recommended that primary care clinicians refer patients to the appropriate psychosocial oncology specialists, mental health professionals, and/or resources in the community.²⁶ After referring to the appropriate resource(s), primary care clinicians should follow-up with patients to check their adherence to recommended therapies and/or assess the need for additional referrals. If a patient has difficulties with adherence to recommendations, primary care clinicians should identify the challenges with adherence and help the patient overcome these obstacles before discussing alternative interventions.¹⁵ The American Psychosocial Oncology Society Web site (apos-society.org/) can help primary care clinicians identify resources for their patients.

Fatigue

Where appropriate, these recommendations build on the recently published ASCO screening, assessment and treatment of cancer-related fatigue guidelines among adult cancer survivors.²⁵

Recommendation 3.6. It is recommended that primary care clinicians (a) should assess for fatigue and treat any causative factors for fatigue, including anemia, thyroid dysfunction, and cardiac dysfunction (LOE = 0); (b) should offer treatment or referral for factors that may impact fatigue (eg, mood disorders, sleep disturbance, pain, and so on) for those who do not have an otherwise identifiable cause of fatigue (LOE = I); and (c) should counsel patients to engage in regular physical activity and refer for cognitive behavioral therapy as appropriate (LOE = I).

Clinical interpretation. Cancer-related fatigue is very common among those treated for cancer, especially those who undergo treatment with radiation therapy and chemotherapy,^{13,25,108} with an estimated prevalence of 28% to 91%.^{109,110} Recommendations for how to screen and assess for fatigue are provided online (see online supporting information) and come from the ASCO guidelines.²⁵ For some, fatigue lasts long after treatment and can significantly interfere with QoL. Treatable causes of fatigue include anemia, thyroid dysfunction, and cardiac dysfunction.²⁵ For those who do not have an identifiable physical cause of fatigue (anemia), contributing factors, such as mood disorders, sleep disturbance, and pain, should be addressed.²⁵ Additional information related to distress/depression and pain can be found under Recommendation 4.5.

A regular exercise regimen can reduce fatigue, help survivors feel better physically and emotionally, and help them cope, as has been demonstrated by several RCTs in breast cancer survivors^{23,25,111} (see also Recommendation 4.3). Cognitive behavioral therapy may also lessen fatigue.^{112,113} There are minimal data to support use of pharmacologic agents for the management of fatigue in this population.²⁵ Interventions should be tailored to the needs and abilities of the individual breast cancer survivor. ASCO has more detailed information on the management of fatigue for cancer survivors (instituteofquality.org/screening-assessment-and-management-fatigue-adult-survivors-cancer-american-society-clinical²⁵).

Bone Health

Recommendation 3.7. It is recommended that primary care clinicians (a) should refer postmenopausal breast cancer survivors for a baseline dual-energy x-ray absorptiometry (DEXA) scan (LOE = 0); and (b) should refer for repeat DEXA scans every 2 years for women who are taking an aromatase inhibitor, premenopausal women who are taking tamoxifen and/or a gonadotropin-releasing hormone (GnRH) agonist, and women who have chemotherapy-induced premature menopause (LOE = 0).

Clinical interpretation. The rate and magnitude of bone loss caused by cancer therapy are significantly higher than normal age-related bone loss.^{114,115} Up to 80% of breast cancer patients experience bone loss.^{116,117} Osteoporosis risk factors unique to patients after cancer therapy include chemotherapy-induced premature menopause, GnRH suppression of gonadal function, anti-estrogen therapies, and glucocorticoids.¹¹⁸ These risk factors are cumulative with other known risk factors, including age, prior fracture history, and family history of fracture.^{119,120} Lifestyle-related factors, including smoking, excess alcohol, inadequate exercise, low calcium, and vitamin D deficiency, are common in this population and increase the risk of osteoporosis.¹¹⁸ Primary care clinicians should manage symptoms as they would in the general population.

Postmenopausal women treated with aromatase inhibitors are at increased risk of osteoporosis and should have initial and periodic (every 2 years) DEXA scan screening. If major risk factors change, then it is reasonable to consider a repeat DEXA scan at 1 year.¹¹⁸ All postmenopausal women or premenopausal women receiving ovarian suppression therapy with GnRH agonists are at risk for developing osteoporosis and should be screened according to the US Preventive Services Task Force³⁰ and the American Association of Clinical Endocrinologists guide for postmenopausal osteoporosis diagnosis and treatment.³¹

Initial strategies to reduce the morbidity associated with bone loss include education about risk factors and a healthy lifestyle. These should include physical activity and regular weight-bearing exercise, avoiding tobacco use, limiting alcohol intake, and consider supplementation with calcium (to achieve a total intake of 1200 mg/d) and vitamin D3 (600-1000 IU/d) for all adults older than 50 years.^{118,121-126} In addition to lifestyle and nutritional interventions, pharmacologic options should be considered in patients at high risk for bone loss and/or fracture.¹¹⁸ Bisphosphonates or denosumab can prevent bone loss and/or treat established osteoporosis.^{118,127-129} However, these drugs do have adverse effects and risks, so that the risk versus benefit of antiresorptive therapy must be carefully considered before starting therapy. Estrogen receptor modulators (raloxifene and tamoxifen) also have antiresorptive properties. However, in one large trial, combining

an SERM and an aromatase inhibitor blunted the reduction in breast cancer recurrence compared with an aromatase inhibitor alone. Therefore, SERMs should not be used for the prevention of osteoporosis in women who are taking an aromatase inhibitor.¹³⁰

Musculoskeletal Health

Recommendation 3.8. It is recommended that primary care clinicians (a) should assess for musculoskeletal symptoms, including pain, by asking patients about their symptoms at each clinical encounter (LOE = 0); and (b) should offer one or more of the following interventions based on clinical indication: acupuncture, physical activity, referral for physical therapy or rehabilitation (LOE = III).

Clinical interpretation. Breast cancer survivors may report difficulties with the ipsilateral upper extremity after surgery, including decreased range of motion, rotator cuff injury, adhesive capsulitis ("frozen shoulder" with stiffness and pain in the shoulder joint), and axillary web syndrome ("cording" in the skin of the inner arm with sensations of pain and tightness that appear as a web or a corded rope).^{131,132} These abnormalities can lead to a decreased ability to perform activities of daily living and can impact employment. Systemic therapies for breast cancer have also been associated with the development of musculoskeletal symptoms.¹³³ The prevalence of musculoskeletal symptoms among breast cancer patients varies greatly: these include limited shoulder range of motion (range, 1.5% to 50% of patients), musculoskeletal pain (range, 12% to 51% of patients), upper limb weakness (range, 18% to 23% of patients), and numbness (range, 29% to 81% of patients).^{131,134-136} In particular, up to 50% of postmenopausal women receiving treatment with aromatase inhibitor medications report arthralgias (joint pain) and myalgias (muscle pain) that are severe enough in 20% of women to lead to treatment discontinuation.^{137,138} These aromatase inhibitor-associated musculoskeletal symptoms are often not responsive to nonsteroidal anti-inflammatory drugs or acetaminophen. Another option for treatment is to change from one antiestrogen therapy to another. Approximately 40% who discontinue the drug may tolerate a different aromatase inhibitor or a different formulation of the aromatase inhibitor. The rest generally tolerate tamoxifen.¹³⁹ Poor compliance/adherence to therapy has been shown to result in an increased risk of breast cancer recurrence, so helping patients manage their symptoms and encouraging drug compliance is important.¹⁴⁰

Physical therapy, including stretching and other exercises, has been shown to be effective for managing postsurgical musculoskeletal symptoms.^{141,142} Recent data from the Hormones and Physical Exercise trial, a prospective cohort study, demonstrated that participation in an intensive exercise regimen resulted in a 20% decrease in aromatase inhibitor-associated pain.¹⁴³ To date, only acupuncture¹⁴⁴ and exercise¹⁴³ have been demonstrated to result in a statistically significant improvement in aromatase inhibitor-associated symptoms.

Pain and Neuropathy

Recommendation 3.9. It is recommended that primary care clinicians (a) should assess for pain and contributing factors for pain with the use of a simple pain scale and comprehensive history of the patient's complaint (LOE = 0); (b) should offer interventions, such as acetaminophen, nonsteroidal anti-inflammatory drugs, physical activity, and/or acupuncture, for pain (LOE = I); (c) should refer to an appropriate specialist, depending on the etiology of the pain once the underlying etiology has been determined (eg, lymphedema specialist,

occupational therapist, and so on; LOE = 0); (d) should assess for peripheral neuropathy and contributing factors for peripheral neuropathy by asking the patient about their symptoms, specifically numbness and tingling in their hands and/or feet, and the characteristics of that symptom (LOE = 0); (e) should offer physical activity for neuropathy (LOE = IA); and (f) should offer duloxetine for patients with neuropathic pain, numbness, and tingling (LOE = IB).

Clinical interpretation. A substantial percentage of breast cancer survivors experience long-term, treatment-related chronic pain that can negatively impact QoL. Published reports demonstrate that from 25% to 60% of breast cancer survivors experience chronic pain as a result of the treatments administered, including surgery, radiation therapy, chemotherapy, and endocrine therapy.^{138,145-148}

Patients should also be evaluated for secondary causes of pain, such as lymphedema or tightness of the chest wall or axilla, and should be referred to lymphedema specialists or occupational therapists as indicated. Once the work-up and assessment are complete and underlying causes are identified or ruled out, chronic pain after breast cancer surgery is typically treated with standard analgesics, including acetaminophen and nonsteroidal anti-inflammatory drugs.

Acupuncture and physical activity have been shown in meta-analyses of RCTs to improve pain among breast cancer survivors and are typically used to complement traditional cancer care. Treatment-related joint pain among breast cancer survivors is quite commonly treated with acupuncture, and many trials and systematic reviews have demonstrated its efficacy in decreasing pain intensity.^{144,149} However, evidence is lacking to show that acupuncture directly benefits breast cancer survivors who are experiencing CIPN. Physical activity has been shown in multiple RCTs to improve pain.¹⁵⁰

Neuropathy, including numbness, tingling, and burning pain, is also common after a diagnosis of breast cancer and subsequent treatment. It is particularly common after surgery and after treatment with taxane-based or platinum-based chemotherapy regimens and is reported in 30% to 40% of patients.¹⁴⁷ ASCO recently published a clinical practice guideline about the prevention and management of CIPN (institutequality.org/prevention-and-management-chemotherapy-induced-peripheral-neuropathy-survivors-adult-cancers),¹⁴ and we endorse that guideline.

Prevention and treatment approaches for the management of CIPN in adult cancer survivors, reproduced from the ASCO guideline, are available online (see online supporting information). In a few small RCTs with breast cancer survivors, physical activity has been shown to improve arthralgias, neuropathy, and neuropathy symptoms.^{143,151} Several pharmaceutical agents have been tested for the management of CIPN. In a randomized, placebo-controlled trial, the serotonin-norepinephrine reuptake inhibitor (SNRI) duloxetine was shown to decrease neuropathic pain significantly more than placebo, and it may also improve numbness and tingling. That trial used an initial dose of 30 mg daily for the first week to reduce the likelihood of nausea; then, the dose was increased to 60 mg daily. The associated relative risk benefit was 30% to 50% in pain reduction.¹⁵² Studies of tricyclic antidepressants and anticonvulsants have not demonstrated consistent significant improvements in symptoms. Additional recommendations for the prevention and management of CIPN can be found in the recently published ASCO guidelines.¹⁴

Infertility

Recommendation 3.10. It is recommended that primary care clinicians should refer survivors of childbearing age who experience

infertility to a specialist in reproductive endocrinology and infertility as soon as possible (LOE = 0).

Clinical interpretation. Infertility as a result of cancer treatment is a potential long-term adverse effect faced by younger breast cancer survivors (younger than age 45 years). When it occurs, it can have a profound impact on a survivor's physical and psychosocial QoL.⁴⁷ Breast cancer patients between ages 20 and 45 years represent 10.9% of all new breast cancer cases in the United States,² making infertility an issue for many younger survivors.¹⁵³

Chemotherapy can be gonadotoxic, leading to reduced fertility or early menopause secondary to premature ovarian failure.^{76,153} Many of the most frequently used chemotherapy agents in the treatment of breast cancer (eg, alkylating agents, platinum agents, and taxanes) are also those that most often lead to premature ovarian failure.¹⁵⁴ The incidence of chemotherapy-related amenorrhea increases with age, as the female ovarian reserve is nonrenewable and diminishes steadily with age. However, there is considerable variation in ovarian reserve among women of similar age.¹⁵⁵ There is limited literature related to the gonadotoxicity of biologics and targeted therapies, and more studies are needed. Primary care clinicians should involve the treating medical oncologist in any potential discussion related to the optimal time for pregnancy after the completion of breast cancer treatment. Premenopausal women who desire pregnancy and are having difficulty conceiving for 6 months or more (or who have had > one miscarriage) should be referred to a fertility specialist. Timely referral is crucial because of the rapid loss of ovarian reserve in these women.

Sexual Health

Recommendation 3.11. It is recommended that primary care clinicians (a) should assess for signs and symptoms of sexual dysfunction or problems with sexual intimacy (LOE = 0); (b) should assess for reversible contributing factors to sexual dysfunction and treat when appropriate (LOE = 0); (c) should offer nonhormonal, water-based lubricants and moisturizers for vaginal dryness (LOE = IA); and (d) should refer for psychoeducational support, group therapy, sexual counseling, marital counseling, or intensive psychotherapy when appropriate (LOE = IA).

Clinical interpretation. Sexual complaints are a common problem among breast cancer survivors that should be assessed. They can include sexual desire disorder/decreased libido (range, 23% to 64% of patients), arousal or lubrication concerns (range, 20% to 48% of patients), orgasmic concerns (range, 16% to 36% of patients), and dyspareunia (range, 35% to 38% of patients).⁷⁶ Patients who receive chemotherapy tend to have more of these sexual concerns than those treated only with surgery and/or radiation.⁴⁹ Treatment with aromatase inhibitors may cause vaginal dryness, dyspareunia (which can be severe), menopausal symptoms, and loss of sexual desire.¹⁵⁶ Radiation therapy can often cause skin fibrosis, loss of sexual sensitivity of the skin, and, uncommonly, cardiac and respiratory damage, all of which negatively impact sexual desire and response.¹⁵⁷

It is important to counsel patients concerning possible sexual dysfunction remedies, including treatments for vaginal dryness. Nonhormonal, water-based lubricants and moisturizers remain the primary treatment.¹⁵⁸ Silicone-based products may last longer than water-based or glycerin-based products. A combination of therapies may provide additional short-term comfort. Hormonal therapies, such as a low-dose estrogen vaginal tablets or an estradiol vaginal ring,

may be recommended for vaginal dryness because of urogenital atrophy, although results commonly take approximately 6 to 12 weeks.^{158,159} The safety of these therapies in women with a history of breast cancer is not well established at this time. The level of estrogen absorption is variable, which raises concerns in patients who have a history of breast cancer. Use of hormonal therapies for women on aromatase inhibitors is not recommended.¹⁶⁰ Treating dyspareunia secondary to vaginal atrophy and stenosis with vaginal dilators or pelvic floor relaxation techniques may be helpful.⁷⁶

Referral for interventions, such as brief psychoeducational support, group therapy, sexual counseling, marital counseling or intensive psychotherapy, should be offered to all breast cancer survivors with sexual complaints specifically addressing possible anxiety, stress, unpleasant symptoms including hot flashes, sexual comfort in lovemaking, and mood changes.^{49,156,161} Taken together, the trio of counseling, over-the-counter treatments, and pharmacologic treatments can do much to ameliorate the sexual issues caused by breast cancer and its management.¹⁵⁶ See Recommendation 3.1 to address body image concerns.

Premature Menopause/Hot Flashes

Recommendation 3.12. It is recommended that primary care clinicians should offer SNRIs, selective serotonin reuptake inhibitors (SSRIs), gabapentin, lifestyle modifications and/or environmental modifications to help mitigate vasomotor symptoms of premature menopausal symptoms (LOE = IA).

Clinical interpretation. Women can experience menopausal symptoms if chemotherapy results in premature cessation of ovarian function or as an adverse effect of endocrine therapies. Vasomotor symptoms are typically more severe in younger survivors because of the abrupt change in hormones¹⁶² and, when present, can have a significant impact on QoL. For younger women on endocrine therapies, 50% to 70% will likely experience hot flashes while on tamoxifen.⁷⁶

Systemic hormone therapy is given rarely, if ever, to patients with breast cancer to control menopausal symptoms. Nonhormonal medications like SNRIs and SSRIs can decrease the intensity and severity of vasomotor symptoms, although they are not approved by the US Food and Drug Administration for this indication. The SNRI venlafaxine has been found to be safe and effective in reducing hot flashes.¹⁶³ There is concern that SSRIs that inhibit the CYP2D6 (CYP450 2D6) enzyme pathway, such as paroxetine, may reduce the conversion of tamoxifen to active metabolites, although a negative impact on breast cancer outcomes has not been conclusively demonstrated.¹⁶⁴ Consistent with recommendations by the ASCO and NCCN, patients should not be screened for CYP2D6.^{9,165} The anticonvulsant gabapentin has also been shown to be effective in reducing hot flashes¹⁶⁶⁻¹⁶⁸ Similarly, the antihypertensive clonidine has been used in clinical practice.¹⁶⁹ Antihypertensive medications and modifications in lifestyle and environment may also help decrease the intensity and severity of menopausal symptoms. In a meta-analysis of RCTs among breast cancer survivors, acupuncture reduced menopausal symptoms and hot flashes.¹⁷⁰ Lifestyle interventions, including rhythmic breathing, vitamins, exercise, and avoiding spicy foods, caffeine, and alcohol, have had variable results. Environmental modifications, such as cool rooms and dressing in layers, can also be helpful. Complementary therapies have been studied, and some have been found to be minimally effective.¹⁷¹

HEALTH PROMOTION

Women with breast cancer are likely to have long-term survival.¹ Thus, enhancing the length and quality of life is an important goal in the care of patients with breast cancer and in creating a survivorship care plan (see Recommendations 4.1 and 5.1 for more information on survivorship care plans). Healthy behaviors are paramount to reducing the risk of second cancers, comorbidities, obesity, and possibly recurrence; improving prognosis; ameliorating cancer-related symptoms¹⁷²⁻¹⁷⁷; and decreasing the risk of mortality.¹⁷⁴⁻¹⁸⁴

Table 7 outlines this information, including physical activity, nutrition, and smoking cessation recommendations for breast cancer survivorship, incorporating existing ACS nutrition and physical activity recommendations²³ (**Table 7: Health Promotion Guidelines**).

Information

Recommendation 4.1. It is recommended that primary care clinicians (a) should assess the information needs of the patient related to breast cancer and its treatment, adverse effects, other health concerns, and available support services (LOE = 0); and (b) should provide or refer survivors to appropriate resources to meet these needs (LOE = 0).

Clinical interpretation. Breast cancer survivors often express unmet needs for information after treatment, including information on the effects of cancer treatment, emotional distress, and lifestyle changes.^{185,186} Younger breast cancer patients can be particularly vulnerable to the physical, emotional, and psychosocial late effects of treatment because of the aggressiveness of their disease, the intensity of the treatment plan,^{47,187} and their younger age, when a cancer diagnosis is not as common or is unexpected.^{188,189} Treatment summaries and individualized survivorship care plans provide survivors with individualized information on their cancer care. However, to date, there is inconsistency in the implementation, comprehensiveness, and perceived helpfulness of such plans for both patients and primary care clinicians.¹⁹⁰⁻¹⁹²

Results from randomized trials of survivorship care plans are mixed,¹⁹³⁻¹⁹⁶ making the direct benefits of survivorship care plans less clear. However, the failure of specialists to provide treatment summaries and survivorship care plans is an obstacle to the ability of primary care clinicians to provide relevant information and care to their patients with a history of breast cancer.

The informational needs of breast cancer survivors and caregivers should be routinely assessed, and information about the long-term and late effects of breast cancer treatment, as well as information on risk reduction and health promotion, should be provided. Resources that may be beneficial to share with patients include the ACS Survivorship Center Web site (cancer.org/survivorshipcenter), the ACS Web site (cancer.org), Journey Forward (journeyforward.org), the ASCO survivor and caregiver site (cancer.net), and the NCCN patient and caregiver resources (nccn.org/patients).

Obesity

Recommendation 4.2. It is recommended that primary care clinicians (a) should counsel survivors to achieve and maintain a healthy weight (LOE = III); and (b) should counsel survivors, if overweight or obese, to limit consumption of high-calorie foods and beverages and increase physical activity to promote and maintain weight loss (LOE = IA, III).

Table 7. Health Promotion Guideline

Recommendation	Level of Evidence ^a
It is recommended that primary care clinicians:	
Recommendation 4.1: Information	
(a) Should assess the information needs of the patient related to breast cancer and its treatment, side effects, other health concerns, and available support services	0 (assessment)
(b) Should provide or refer survivors to appropriate resources to meet these needs	0 (referral)
Recommendation 4.2: Obesity	
(a) Should counsel survivors to achieve and maintain a healthy weight	0 (maintenance)
(b) Should counsel survivors if overweight or obese to limit consumption of high-calorie foods and beverages and increase physical activity to promote and maintain weight loss	IA, III (weight loss)
Recommendation 4.3: Physical activity	
Should counsel survivors to engage in regular physical activity consistent with the ACS guideline, and specifically	
(a) Should avoid inactivity and return to normal daily activities as soon as possible following diagnosis	III (avoid inactivity)
(b) Should aim for at least 150 min of moderate or 75 min of vigorous aerobic exercise per wk	I, IA (aerobic exercise)
(c) Should include strength training exercises at least 2 d per wk; emphasize strength training for women treated with adjuvant chemotherapy or hormone therapy	IA (strength training)
Recommendation 4.4: Nutrition	
Should counsel survivors to achieve a dietary pattern that is high in vegetables, fruits, whole grains, and legumes; low in saturated fats; and limited in alcohol consumption	IA, III (nutrition); 0 (alcohol)
Recommendation 4.5: Smoking cessation	
Should counsel survivors to avoid smoking and refer survivors who smoke to cessation counseling and resources	I

Abbreviation: ACS, American Cancer Society.

^aLevel of evidence: I indicates meta-analyses of randomized controlled trials (RCTs); IA, RCT of breast cancer survivors; IB, RCT based on cancer survivors across multiple sites; IC, RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing fatigue, lymphedema, etc); IIA, nonrandomized controlled trial (non-RCT) based on breast cancer survivors; IIB, non-RCT based on cancer survivors across multiple sites; IIC, non-RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing urinary incontinence, erectile dysfunction, etc); III, case-control or prospective cohort study; 0, expert opinion, observation, clinical practice, literature review, or pilot study.

Clinical interpretation. Approximately 62% of breast cancer survivors are overweight/obese (have a body mass index of least 25 kg/m²), of whom 30% are classified as obese (body mass index, 30 kg/m² or higher).¹⁹⁷ Sufficient evidence documents obesity as a risk factor for postoperative complications, second cancers, risk of recurrence, development of diabetes, and other issues. Conversely, weight loss mitigates symptoms and improves QoL.¹⁹⁸ ASCO issued a position statement underscoring the need for oncology providers to counsel their patients about achieving a healthy weight.¹⁹⁹ Primary care clinicians should also counsel cancer survivors to achieve or maintain a healthy weight¹⁹⁸ and refer to multicomponent obesity treatment programs where appropriate.²⁰⁰

Physical Activity

Recommendation 4.3. It is recommended that primary care clinicians should counsel survivors to engage in regular physical activity consistent with the ACS guideline²³ and, specifically: (a) should avoid inactivity and return to normal daily activities as soon as possible after diagnosis (LOE = III), (b) should aim for at least 150 minutes of moderate or 75 minutes of vigorous aerobic exercise per week (LOE = I, IA), and (c) should include strength training exercises at least 2 days per week. Strength training should be emphasized for women who are treated with adjuvant chemotherapy or hormone therapy (LOE = IA).

Clinical interpretation. Approximately 32% of cancer survivors meet the recommendations for physical activity.²⁰¹ Numerous systematic reviews and meta-analyses have documented the many health benefits from physical activity in breast cancer survivors, including mitigating treatment adverse effects (ie, fatigue) and improving QoL and physical functioning. The data suggesting a potential survival benefit of physical activity come from observational data, with the

most recent review of this literature citing 16 studies that reported an average relative risk of 0.72 for breast cancer deaths in physically active breast cancer survivors (95% CI, 0.60-0.85) and 0.52 for all-cause mortality (95% CI, 0.42-0.64).²⁰² Breast cancer survivors should be advised to return to normal daily activities as soon as possible after diagnosis and to continue engaging in regular physical activity. Breast cancer survivors should strive to exercise at least 150 minutes moderately or 75 minutes vigorously per week and should include strength training exercises at least 2 days per week.²³ Additional details regarding the amount of time needed for each strength training session is less clear. However, studies indicate that from 37% to 53% of breast cancer survivors meet the aerobic guideline,^{203,204} and 23% meet the strength training guideline.²⁰⁵ Observational evidence suggests greater amounts of activity may be needed, although the evidence is insufficient to make it a recommendation at this time; aerobic exercise of 3 hours or more per week may be needed to improve breast cancer survival.^{184,206}

Nutrition

Recommendation 4.4. It is recommended that primary care clinicians should counsel survivors to achieve a dietary pattern that is high in vegetables, fruits, whole grains, and legumes; low in saturated fats (LOE = IA, III); and limited in alcohol consumption (LOE = 0).

Clinical interpretation. Only 18% to 34% of breast cancer survivors report eating five or more fruit and vegetables daily.²⁰⁴ Eating a diet characterized by high amounts of vegetables, fruits, whole grains, and legumes (v a typical Western diet) has been associated with a reduced risk (range, 15% to 43%) in all-cause mortality.²⁰⁷⁻²⁰⁹ Data from the two large RCTs of diet interventions in breast cancer survivors suggest that dietary change sufficient to result in weight loss may

be needed to favorably impact breast cancer recurrence and prognosis.^{210,211} According to the ACS Nutrition and Physical Activity Guidelines, alcohol consumption should be limited to no more than one drink per day for women, as in the general population.^{212,213} Data are inconsistent but suggest that breast cancer survivors who consume more than three to four drinks per week are at increased risk for breast cancer recurrence.²¹⁴

Studies have shown that the carcinogenic ingredients of alcohol increase the risk of developing many types of cancers²¹⁵⁻²¹⁷ when alcohol, regardless of the type (ie, wine, beer, and so on), is consumed in excess of daily recommended limits.²¹⁸ Approximately 7% of breast cancer survivors report excessive drinking.²⁰³ Based on these data, survivors should be counseled to: achieve a dietary pattern that is high in vegetables, fruits, whole grains, and legumes; limit alcohol intake to no more than one drink per day; and follow the ACS Guidelines on Nutrition and Physical Activity for Cancer Survivors²³ with a focus on successful weight management.

Smoking Cessation

Recommendation 4.5. It is recommended that primary care clinicians should counsel survivors to avoid smoking and refer survivors who smoke to cessation counseling and resources (LOE = I).

Clinical interpretation. Approximately 10% to 12% of breast cancer survivors smoke.^{203,204} Numerous observational studies show that women who smoke at the time of diagnosis have substantially worse breast cancer–specific and overall survival than former and never-smokers.²¹⁹ A recent meta-analysis of this observational work documents a 33% increased risk of mortality from breast cancer in women who are smokers at diagnosis compared with former smokers.²¹⁹ Survivors should discontinue smoking and avoid subsequent tobacco product use. Clinicians should identify smokers and motivate and encourage patients to quit through cessation programs, brochures and pamphlets, counseling, pharmacotherapy, and regular follow-up.²²⁰ These programs should be initiated at the time of initial diagnosis if possible.

CARE COORDINATION/PRACTICE IMPLICATIONS

There are no clear guidelines for the shared care and comanagement of patients with breast cancer after the completion of active treatment (Table 8). The time to transition is variable and depends

on medical, geographic, and resource restraints. Some patients return to their primary care clinician immediately after treatment is completed.²²¹⁻²²³ Others may choose to transition to their follow-up care only after they are considered at little or low risk for disease recurrence or late effects of cancer treatment.^{224,225} Several RCTs show that care led by primary care clinicians is as effective as hospital-led or specialist-led care, including similar rates of recurrence-related, serious clinical events; levels of health-related QoL²²⁶; and patient satisfaction with care.²²⁷

Breast cancer survivors may continue to see their oncology team for follow-up disease surveillance; however, they should also be seen by their primary care clinician for health maintenance and management of comorbidities that may or may not be related to their cancer diagnosis and treatment.^{222,228}

Survivorship Care Plan

Recommendation 5.1. It is recommended that primary care clinicians should consult with the cancer treatment team and obtain a treatment summary and survivorship care plan (LOE = 0, III).

Clinical interpretation. Survivorship care plans are recommended as an important tool to facilitate communication and allocation of responsibility during the transition from active treatment to survivorship care.^{4,229} A summary of a patient's diagnosis and treatment received should be provided by the oncology care team when a patient with breast cancer transitions care to other providers; a treatment summary should describe the type and stage/side of the cancer, type of surgery, the name of the chemotherapy/hormones/biologics and cumulative doses of chemotherapy, and the types and cumulative doses of radiation therapy, including the fields and extent of the radiation.^{4,190} Patients can initiate the building of a survivorship care plan process on the ASCO Web site (cancer.net/survivorship/follow-care-after-cancer-treatment/asco-cancer-treatment-and-survivorship-care-plans; at journeyforward.org/ or livestrongcareplan.org/).

Ideally, the oncology team should also work with the patient to develop an individualized cancer survivorship care plan for breast cancer survivors. This care plan guides recommendations for the type and timing of follow-up imaging, laboratory tests, and office visits. The care plan should include information on the risk for late effects of treatment and what to watch for specifically based on the type of cancer and treatment received. Survivors should be assessed for the

Table 8. Care Coordination Guideline

Recommendation	Level of Evidence ^a
It is recommended that primary care clinicians:	
Recommendation 5.1: Survivorship care plan	
Should consult with the cancer treatment team and obtain a treatment summary and survivorship care plan	0, III
Recommendation 5.2: Communication with oncology team	
Should maintain communication with the oncology team throughout your patient's diagnosis, treatment, and posttreatment care to ensure care is evidence-based and well-coordinated	0
Recommendation 5.3: Inclusion of family	
Should encourage the inclusion of caregivers, spouses, or partners in usual breast cancer survivorship care and support	0

^aLevel of evidence: I indicates meta-analyses of randomized controlled trials (RCTs); IA, RCT of breast cancer survivors; IB, RCT based on cancer survivors across multiple sites; IC, RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing fatigue, lymphedema, etc); IIA, nonrandomized controlled trial (non-RCT) based on breast cancer survivors; IIB, non-RCT based on cancer survivors across multiple sites; IIC, non-RCT not based on cancer survivors but on general population experiencing a specific long-term or late effect (eg, managing urinary incontinence, erectile dysfunction, etc); III, case-control or prospective cohort study; 0, expert opinion, observation, clinical practice, literature review, or pilot study.

presence of these physical effects (eg, cardiovascular issues, musculoskeletal issues) as well as psychosocial effects (eg, cognitive dysfunction, depression, FOR, body image, and sexual dysfunction) and should be referred to the appropriate providers and services.

However, the field of oncology is broadly struggling with how to best meet this recommendation and in identifying the specific benefits of such care plans. Various tools and strategies to facilitate the creation and distribution of these care plans are being actively considered for all tumor sites, including breast cancer. Currently, challenges in workflow and tools make this difficult, but the field is working toward a sustainable solution.

Communication With Oncology Team

Recommendation 5.2. It is recommended that primary care clinicians should maintain communication with the oncology team throughout your patient's diagnosis, treatment, and post-treatment care to ensure care is evidence based and well coordinated (LOE = 0).

Clinical interpretation. Communication and cooperation among providers and survivors are critical, with the oncology team providing concrete recommendations for care when needed or requested by other providers.²²⁹ Clear communication regarding the respective roles of different members of the health care team is critical to a successful transition to survivorship care.

The primary care clinician should serve as a general medical care coordinator throughout the spectrum of breast cancer detection and aftercare, focusing on evidence-based preventive care and the management of preexisting comorbid conditions; regularly addressing the patient's overall physical and psychosocial status; making appropriate referrals for psychosocial, rehabilitative, or other specialist care as needed; and coordinating those components of survivorship care that are agreed on with the treating clinicians. Treatment of breast cancer is complex and rapidly changing; therefore, decisions about and coordination of cancer treatment should be left to the oncology team.

Inclusion of Family

Recommendation 5.3. It is recommended that primary care clinicians should encourage the inclusion of caregivers, spouses, or partners in usual breast cancer survivorship care and support (LOE = 0).

Clinical interpretation. Caregivers have to cope with the physical aftermath of the survivors' treatment and manage their long-term and late effects in addition to their own psychosocial and physical unmet needs.²³⁰ Research has shown that, for 14 to 24 months after a survivor's cancer diagnosis, caregivers provide consistent, continuing care involvement²³¹ for patients after breast cancer treatment. Successful coordination of care involves not only a comprehensive care team, including primary care clinicians, but also the informal caregivers (usually the spouse/partner/family member) who provide ongoing care to cancer survivors in the home.²³² Furthermore, most caregivers are older adults who are also managing their own health problems. When possible, primary care clinicians should include caregivers of breast cancer survivors in all follow-up care appointments to optimize survivor wellness.

DISCUSSION

A significant limitation of this guideline is the limited evidence base with which to provide clear and specific recommendations for the

prevention and management of long-term and late effects of cancer survivors. There are few prospective, RCTs testing interventions among breast cancer survivors, although studies in breast cancer survivors dominate the survivorship literature. The majority of the citations characterizing the risk and magnitude of risk of late effects and management recommendations relied predominantly on case-control studies with fewer than 500 participants and reviews that combined studies with various outcome measures. There were several cohort studies that used population-based data to estimate the risk of late effects.

Another limitation is the reliance on previous guidelines for surveillance and symptom management. In addition, the literature review was not managed by a clinical epidemiologist because of limited resources and, instead, was conducted by project staff and an ACS librarian. Furthermore, the guidelines did not result directly from the development of specific clinical questions asked before the literature review; and guidelines included in the literature review were not evaluated through an instrument such as the Rigor of Development subscale of the Appraisal for Guidelines for Research and Evaluation (AGREE II). Recommendations are based on current evidence in the literature, but most evidence is not sufficient to warrant a strong recommendation. Rather, recommendations should be largely seen as possible management strategies given the current limited evidence base.

HEALTH DISPARITIES

Although the ACS/ASCO clinical practice guidelines represent expert recommendations on the best practices in disease management to provide the highest level of cancer care, it is important to note that many patients have limited access to medical care. Racial and ethnic disparities in health care contribute significantly to this problem in the United States. Patients with cancer who are members of racial/ethnic minorities suffer disproportionately from comorbidities, experience more substantial obstacles to receiving care, are more likely to be uninsured, and are at greater risk of receiving poor-quality care than other Americans.²³³⁻²³⁶ Many other patients lack access to care because of their geographic location and distance from appropriate treatment facilities. Awareness of these disparities in access to care should be considered in the context of this clinical practice guideline, and health care providers should strive to deliver the highest level of cancer care to these vulnerable populations.

The ACS and ASCO believe that cancer clinical trials are vital to inform medical decisions and improve cancer care and that all patients should have the opportunity to participate.

MULTIPLE CHRONIC CONDITIONS

Creating evidence-based recommendations to inform the treatment of patients with additional chronic conditions, a situation in which the patient may have two or more such conditions—referred to as multiple chronic conditions (MCCs)—is challenging.²³⁷ Patients with MCCs are a complex and heterogeneous population, making it difficult to account for all of the possible permutations to develop specific recommendations for care. In addition, the best available evidence for treating index conditions, such as cancer, is often from clinical trials in which study selection criteria may exclude these patients to avoid potential interaction effects or confounding of results associated with MCCs. As a result, the reliability of outcome data from these studies may be limited, thereby creating constraints for expert

groups to make recommendations for care in this heterogeneous patient population.

Because many patients for whom guideline recommendations apply present with MCCs, any treatment plan needs to take into account the complexity and uncertainty created by the presence of MCCs and should highlight the importance of shared decision making regarding guideline use and implementation. Therefore, in consideration of recommended care for the target index condition, clinicians should review all other chronic conditions present in the patient and take those conditions into account when formulating the treatment and follow-up plan.

In light of these considerations, practice guidelines should provide information on how to apply the recommendations for patients with MCCs, perhaps as a qualifying statement for recommended care. This may mean that some or all of the recommended care options are modified or not applied, as determined by best practice in consideration of any MCCs.

EXTERNAL REVIEW

After finalization by the workgroup, the guideline was sent to additional internal experts (ACS and ASCO) and external experts for review and comment before submission for publication. Review comments were reviewed by the expert panel and integrated into the final article before approval by the Clinical Practice Guideline Committee.

GUIDELINE IMPLEMENTATION

ACS/ASCO guidelines are developed for implementation across health settings. Barriers to implementation include the need to increase awareness of the guideline recommendations among front-line practitioners and survivors of cancer and caregivers and also to provide adequate services in the face of limited resources. The guideline Bottom Line Box (Table 1) was designed to facilitate the implementation of recommendations. This guideline will be distributed widely through the ASCO Practice Guideline Implementation Network. ACS/ASCO guidelines are posted on the ACS and ASCO websites. This joint ACS-ASCO guideline is published in both *CA: A Cancer Journal for Clinicians* and in *Journal of Clinical Oncology*.

SUMMARY

Breast cancer survivors face potentially significant impacts of cancer and its treatment and deserve high-quality, comprehensive, coordinated clinical follow-up care. Primary care clinicians must consider each patient's individual risk profile and preferences of care to address physical and psychosocial impacts. Survivors should be provided support to address FOR, depression, anxiety, cognitive

impairment, body image issues, sexual concerns, functional changes and physical impairments, relationship changes, other social role difficulties, employment concerns, and financial challenges, among others. Breast cancer survivors also need to be counseled on health promotion strategies to minimize and mitigate long-term and late effects, to ameliorate comorbid health conditions, and to potentially increase survival.

To clarify the roles of all clinicians working with cancer survivors, we concur with the Institute of Medicine that survivors and primary care clinicians should receive a survivorship care plan that includes a concise summary of treatment as well as a clinical follow-up care plan. Ideally, this plan would be constructed in partnership with the survivor to identify and prioritize goals for survivorship care and would be communicated to the patient to ensure their understanding of individual risks; recommended tests, procedures, and supportive care strategies; and how to optimize wellness. Survivorship care should be coordinated with treating cancer specialists.

ADDITIONAL RESOURCES

In addition to this guideline, tools and resources are available to assist primary care clinicians in implementing these recommendations. *CA* offers a Patient Page (onlinelibrary.wiley.com/doi/10.3322/caac.21322/pdf) to help patients understand how to use this guideline to talk to their doctor about surveillance and screening, symptom management, healthy behaviors, and care coordination. *CA* also offers free continuing medical education and free continuing nursing education for this article at acsjournals.com/ce as an additional resource for physicians and nurses. The Survivorship Center also offers The GW Cancer Institute's Cancer Survivorship E-Learning Series for Primary Care Providers (The E-Learning Series), a free, innovative, online, continuing education program to educate primary care clinicians about how to better understand and care for survivors in the primary care setting. Continuing education credits are available at no cost to physicians, nurse practitioners, nurses, and physician assistants for each 1-hour module. Learn more about The E-learning Series at cancersurvivorshipcentereducation.org. Visit asco.org/guidelineswiki to provide comments on the guideline or to submit new evidence.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at www.jco.org.

AUTHOR CONTRIBUTIONS

Manuscript writing: All authors

Final approval of manuscript: All authors

REFERENCES

1. DeSantis CE, Lin CC, Mariotto AB, et al: Cancer treatment and survivorship statistics, 2014. *CA Cancer J Clin* 64:252-271, 2014
2. Howlader N, Noone AM, Krapcho M, et al, Eds. SEER Cancer Statistics Review, 1975-2012 [seer.cancer.gov/csr.1975_2012/, based on the November 2014 SEER data submission, posted to the SEER website April 2015]. Bethesda, MD National Cancer Institute, 2015
3. Ganz PA: Survivorship: Adult cancer survivors. *Prim Care* 36:721-741, 2009
4. Institute of Medicine and National Research Council of the National Academies: From cancer patient to cancer survivor: Lost in transition. Committee on Cancer Survivorship: Improving Care and

Quality of Life, National Cancer Policy Board. Washington, DC, The National Academies Press, 2005

5. Institute of Medicine (US) Committee on Psychosocial Services to Cancer Patients/Families in a Community Setting: Cancer Care for the Whole Patient: Meeting Psychosocial Health Needs. Washington, DC, The National Academies Press, 2008

6. Earle CC, Ganz PA: Cancer survivorship care: Don't let the perfect be the enemy of the good. *J Clin Oncol* 30:3764-3768, 2012

7. Ganz PA, Earle CC, Goodwin PJ: Journal of Clinical Oncology update on progress in cancer survivorship care and research. *J Clin Oncol* 30:3655-3656, 2012

8. Cowens-Alvarado R, Sharpe K, Pratt-Chapman M, et al: Advancing survivorship care through the National Cancer Survivorship Resource Center: Developing American Cancer Society guidelines for primary care providers. *CA Cancer J Clin* 63:147-150, 2013

9. Gradishar WJ, et al: NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) Breast Cancer Version 2.2015. ©2015 National Comprehensive Cancer Network, Inc. NCCN.org. Accessed April 1, 2015

10. Denlinger CS, et al: NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) Survivorship Version 1.2015. © 2015 National Comprehensive Cancer Network, Inc. NCCN.org. Accessed April 1, 2015

11. Khatcheressian JL, Hurley P, Bantug E, et al: Breast cancer follow-up and management after primary treatment: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol* 31:961-965, 2013

12. American Society of Clinical Oncology: Cancer Survivorship. asco.org/practice-research/cancer-survivorship. Accessed May 15, 2015

13. Bower JE: Cancer-related fatigue—mechanisms, risk factors, and treatments. *Nat Rev Clin Oncol* 11:597-609, 2014

14. Hershman DL, Lacchetti C, Dworkin RH, et al: Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol* 32:1941-1967, 2014

15. Andersen BL, DeRubeis RJ, Berman BS, et al: Screening, assessment, and care of anxiety and depressive symptoms in adults with cancer: An American Society of Clinical Oncology guideline adaptation. *J Clin Oncol* 32:1605-1619, 2014

16. Skolarus TA, Wolf AM, Erb NL, et al: American Cancer Society prostate cancer survivorship care guidelines. *CA Cancer J Clin* 64:225-249, 2014

17. Siegel RL, Miller KD, Jemal A: Cancer statistics, 2015. *CA Cancer J Clin* 65:5-29, 2015

18. Jatoti I: Options in breast cancer local therapy: Who gets what? *World J Surg* 36:1498-1502, 2012

19. Jatoti I, Proschan MA: Randomized trials of breast-conserving therapy versus mastectomy for primary breast cancer: A pooled analysis of updated results. *Am J Clin Oncol* 28:289-294, 2005

20. Kerrigan D, Waters P, Ryan M, et al: Follow-up arrangements for breast cancer patients; is it appropriate to transfer surveillance to general practitioners? *Ir Med J* 107:273-275, 2014

21. Saslow D, Boetes C, Burke W, et al: American Cancer Society guidelines for breast screening with MRI as an adjunct to mammography. *CA Cancer J Clin* 57:75-89, 2007

22. Moyer VA: Risk assessment, genetic counseling, and genetic testing for BRCA-related cancer in

women: US Preventive Services Task Force recommendation statement. *Ann Intern Med* 160:271-281, 2014

23. Rock CL, Doyle C, Demark-Wahnefried W, et al: Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin* 62:242-274, 2012

24. Brawley O, Byers T, Chen A, et al: New American Cancer Society process for creating trustworthy cancer screening guidelines. *JAMA* 306:2495-2499, 2011

25. Bower JE, Bak K, Berger A, et al: Screening, assessment, and management of fatigue in adult survivors of cancer: An American Society of Clinical Oncology clinical practice guideline adaptation. *J Clin Oncol* 32:1840-1850, 2014

26. Holland JC, et al: NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) Distress Management Version 2.2015. © 2015 National Comprehensive Cancer Network, Inc. NCCN.org. Accessed April 1, 2015

27. Loren AW, Mangu PB, Beck LN, et al: Fertility preservation for patients with cancer: American Society of Clinical Oncology clinical practice guideline update. *J Clin Oncol* 31:2500-2510, 2013

28. The University of Texas MD Anderson Cancer Center: Survivorship—Invasive Breast Cancer. mdanderson.org/education-and-research/resources-for-professionals/clinical-tools-and-resources/practice-algorithms/survivorship-breast-invasive-web-algorithm.pdf Accessed April 7, 2014

29. Helfand M, Carson S: US Preventive Services Task Force Evidence Syntheses, formerly Systematic Evidence Reviews. Screening for Lipid Disorders in Adults: Selective Update of 2001 US Preventive Services Task Force Review. Rockville, MD: Agency for Healthcare Research and Quality; 2008

30. US Preventive Services Task Force: Screening for osteoporosis: US Preventive Services Task Force recommendation statement. *Ann Intern Med* 154:356-364, 2011

31. Watts NB, Bilezikian JP, Camacho PM, et al: American Association of Clinical Endocrinologists Medical Guidelines for Clinical Practice for the diagnosis and treatment of postmenopausal osteoporosis. *Endocr Pract* 16(suppl 3):1-37, 2010

32. Chalasani P, Downey L, Stopeck AT: Caring for the breast cancer survivor: A guide for primary care physicians. *Am J Med* 123:489-495, 2010

33. Harris SR, Schmitz KH, Campbell KL, et al: Clinical practice guidelines for breast cancer rehabilitation: Syntheses of guideline recommendations and qualitative appraisals. *Cancer* 118:2312-2324, 2012

34. Stout NL, Binkley JM, Schmitz KH, et al: A prospective surveillance model for rehabilitation for women with breast cancer. *Cancer* 118:2191-2200, 2012

35. van Londen G, Beckjord E, Dew M, et al: Breast cancer survivorship symptom management: Current perspective and future development. *Breast Cancer Manag* 2:71-81, 2013

36. Ganz PA, Yip CH, Gralow JR, et al: Supportive care after curative treatment for breast cancer (survivorship care): resource allocations in low- and middle-income countries. A Breast Health Global Initiative 2013 consensus statement. *Breast* 22:606-615, 2013

37. Centers for Disease Control and Prevention Vaccines: VPD-VAC/Who Should NOT Get Vaccinated? cdc.gov/vaccines/vpd-vac/should-not-vacc.htm. Accessed August 17, 2015

38. Rojas MP, Telaro E, Russo A, et al: Follow-up strategies for women treated for early breast cancer

[serial online]. *Cochrane Database Syst Rev* (1):CD001768, 2005

39. Salani R, Andersen BL: Gynecologic care for breast cancer survivors: Assisting in the transition to wellness. *Am J Obstet Gynecol* 206:390-397, 2012

40. Del Turco M, Palli D, Cariddi A, et al: Intensive diagnostic follow-up after treatment of primary breast cancer: A randomized trial. *JAMA* 271:1593-1597, 1994

41. Impact of follow-up testing on survival and health-related quality of life in breast cancer patients. A multicenter randomized controlled trial. The GIVIO Investigators. *JAMA* 271:1587-1592, 1994

42. Henry LN, Hayes DF, Ramsey SD, et al: Promoting quality and evidence-based care in early-stage breast cancer follow-up [serial online]. *J Natl Cancer Inst* 2014;106:dju034

43. Ruddy K, Mayer E, Partridge A: Patient adherence and persistence with oral anti-cancer treatment. *CA Cancer J Clin* 59:56-66, 2009

44. Smith RA, Manassaram-Baptiste D, Brooks D, et al: Cancer screening in the United States, 2015: a review of current American Cancer Society guidelines and current issues in cancer screening. *CA Cancer J Clin* 65:30-54, 2015

45. Runowicz CD: Gynecologic surveillance of women on tamoxifen: First do no harm. *J Clin Oncol* 18:3457-3458, 2000

46. Falk Dahl CA, Reinertsen KV, Nesvold IL, et al: A study of body image in long-term breast cancer survivors. *Cancer* 116:3549-3557, 2010

47. Partridge AH: Cancer survivorship and the young breast cancer patient: Addressing the important issues. *Oncologist* 18:e19-e20, 2013

48. Avis NE, Crawford S, Manuel J: Quality of life among younger women with breast cancer. *J Clin Oncol* 23:3322-3330, 2005

49. Fobair P, Spiegel D: Concerns about sexuality after breast cancer. *Cancer J* 15:19-26, 2009

50. Rosenberg SM, Tamimi RM, Gelber S, et al: Body image in recently diagnosed young women with early breast cancer. *Psychooncology* 22:1849-1855, 2013

51. Delanian S, Porcher R, Balla-Mekias S, et al: Randomized, placebo-controlled trial of combined pentoxifylline and tocopherol for regression of superficial radiation-induced fibrosis. *J Clin Oncol* 21:2545-2550, 2003

52. Haddad P, Kalaghchi B, Amouzegar-Hashemi F: Pentoxifylline and vitamin E combination for superficial radiation-induced fibrosis: A phase II clinical trial. *Radiother Oncol* 77:324-326, 2005

53. Magnusson M, Hoglund P, Johansson K, et al: Pentoxifylline and vitamin E treatment for prevention of radiation-induced side-effects in women with breast cancer: A phase two, double-blind, placebo-controlled randomised clinical trial (Ptx-5). *Eur J Cancer* 45:2488-2495, 2009

54. Okunieff P, Augustine E, Hicks JE, et al: Pentoxifylline in the treatment of radiation-induced fibrosis. *J Clin Oncol* 22:2207-2213, 2004

55. Scott JL, Kayser K: A review of couple-based interventions for enhancing women's sexual adjustment and body image after cancer. *Cancer J* 15:48-56, 2009

56. Armer JM: The problem of post-breast cancer lymphedema: Impact and measurement issues. *Cancer Invest* 23:76-83, 2005

57. Armer JM, Stewart BR: Post-breast cancer lymphedema: Incidence increases from 12 to 30 to 60 months. *Lymphology* 43:118-127, 2010

58. Radina ME, Armer JM, Stewart BR: Making self-care a priority for women at risk of breast cancer-related lymphedema. *J Fam Nurs* 20:226-249, 2014

59. Krag DN, Anderson SJ, Julian TB, et al: Sentinel-lymph-node resection compared with conventional axillary-lymph-node dissection in clinically node-negative patients with breast cancer: Overall survival findings from the NSABP B-32 randomised phase 3 trial. *Lancet Oncol* 11: 927-933, 2010
60. Warren LE, Miller CL, Horick N, et al: The impact of radiation therapy on the risk of lymphedema after treatment for breast cancer: A prospective cohort study. *Int J Radiat Oncol Biol Phys* 88:565-571, 2014
61. Fu MR, Rosedale M: Breast cancer survivors' experiences of lymphedema-related symptoms. *J Pain Symptom Manage* 38:849-859, 2009
62. International Society of Lymphology. The diagnosis and treatment of peripheral lymphedema: 2013 consensus document of the international society of lymphology. *Lymphology* 46:1-11, 2013
63. Shaitelman SF, Cromwell KD, Rasmussen JC, et al: Recent progress in the treatment and prevention of cancer-related lymphedema. *CA Cancer J Clin* 65:55-81, 2015
64. Ostby PL, Armer JM, Dale PS, et al: Surveillance recommendations in reducing risk of and optimally managing breast cancer-related lymphedema. *J Pers Med* 4:424-447, 2014
65. Coriddi M, Khansa I, Stephens J, et al: Analysis of factors contributing to severity of breast cancer-related lymphedema. *Ann Plast Surg* 74:22-25, 2015
66. Schmitz KH, Ahmed RL, Troxel AB, et al: Weight lifting for women at risk for breast cancer-related lymphedema: A randomized trial. *JAMA* 304: 2699-2705, 2010
67. Cemal Y, Pusic A, Mehrara BJ: Preventative measures for lymphedema: Separating fact from fiction. *J Am Coll Surg* 213:543-551, 2011
68. Ahmed RL, Thomas W, Yee D, et al: Randomized controlled trial of weight training and lymphedema in breast cancer survivors. *J Clin Oncol* 24:2765-2772, 2006
69. Dominick SA, Natarajan L, Pierce JP, et al: Patient compliance with a health care provider referral for an occupational therapy lymphedema consult. *Support Care Cancer* 22:1781-1787, 2014
70. Darby SC, Ewertz M, McGale P, et al: Risk of ischemic heart disease in women after radiotherapy for breast cancer. *N Engl J Med* 368:987-998, 2013
71. Barton M: Cholesterol and atherosclerosis: Modulation by oestrogen. *Curr Opin Lipidol* 24: 214-220, 2013
72. Chalasani P, Livingston R: Differential chemotherapeutic sensitivity for breast tumors with "BRCAness": a review. *Oncologist* 18:909-916, 2013
73. Curigliano G, Cardinale D, Suter T, et al: Cardiovascular toxicity induced by chemotherapy, targeted agents and radiotherapy: ESMO Clinical Practice Guidelines. *Ann Oncol* 23:vii155-vii166, 2012
74. Bell LN, Nguyen AT, Li L, et al: Comparison of changes in the lipid profile of postmenopausal women with early stage breast cancer treated with exemestane or letrozole. *J Clin Pharmacol* 52: 1852-1860, 2012
75. Mouridsen H, Giobbie-Hurder A, Goldhirsch A, et al: Letrozole therapy alone or in sequence with tamoxifen in women with breast cancer. *N Engl J Med* 361:766-776, 2009
76. Stan D, Loprinzi CL, Ruddy KJ. Breast cancer survivorship issues. *Hematol Oncol Clin North Am* 27:805-827, ix, 2013
77. Von Ah D, Habermann B, Carpenter JS, et al: cancer survivors. *Eur J Oncol Nurs* 17:236-241, 2013
78. Janelsins MC, Kohli S, Mohile SG, et al: An update on cancer- and chemotherapy-related cognitive dysfunction: Current status. *Semin Oncol* 38: 431-438, 2011
79. Reid-Arndt SA, Yee A, Perry MC, et al: Cognitive and psychological factors associated with early posttreatment functional outcomes in breast cancer survivors. *J Psychosoc Oncol* 27:415-434, 2009
80. Reuter-Lorenz P, Cimprich B: Cognitive function and breast cancer: Promise and potential insights from functional brain imaging. *Breast Cancer Res Treat* 137:33-43, 2013
81. Bruno J, Hosseini SM, Kesler S: Altered resting state functional brain network topology in chemotherapy-treated breast cancer survivors. *Neurobiol Dis* 48:329-338, 2012
82. Dumas JA, Makarewicz J, Schaubhut GJ, et al: Chemotherapy altered brain functional connectivity in women with breast cancer: A pilot study. *Brain Imaging Behav* 7:524-532, 2013
83. Falletti MG, Sanfilippo A, Maruff P, et al: The nature and severity of cognitive impairment associated with adjuvant chemotherapy in women with breast cancer: A meta-analysis of the current literature. *Brain Cogn* 59:60-70, 2005
84. Hurria A, Lachs M: Is cognitive dysfunction a complication of adjuvant chemo-therapy in the older patient with breast cancer? *Breast Cancer Res Treat* 103:259-268, 2007
85. Ganz PA, Petersen L, Castellon SA, et al: Cognitive function after the initiation of adjuvant endocrine therapy in early-stage breast cancer: An observational cohort study. *J Clin Oncol* 32: 3559-3567, 2014
86. Ahles TA, Saykin AJ, McDonald BC, et al: Longitudinal assessment of cognitive changes associated with adjuvant treatment for breast cancer: Impact of age and cognitive reserve. *J Clin Oncol* 28: 4434-4440, 2010
87. Ahles TA, Saykin AJ: Breast cancer chemotherapy-related cognitive dysfunction. *Clin Breast Cancer* 3(suppl 3):S84-S90, 2002
88. Small BJ, Rawson KS, Walsh E, et al: Catechol-O-methyltransferase genotype modulates cancer treatment-related cognitive deficits in breast cancer survivors. *Cancer* 117:1369-1376, 2011
89. Kohli S, Fisher SG, Tra Y, et al: The effect of modafinil on cognitive function in breast cancer survivors. *Cancer* 115:2605-2616, 2009
90. Von Ah D, Jansen C, Allen DH, et al: Putting evidence into practice: Evidence-based interventions for cancer and cancer treatment-related cognitive impairment. *Clin J Oncol Nurs* 15: 607-615, 2011
91. Von Ah D, Carpenter JS, Saykin A, et al: Advanced cognitive training for breast cancer survivors: A randomized controlled trial. *Breast Cancer Res Treat* 135:799-809, 2012
92. Ercoli LM, Petersen L, Hunter AM, et al: Cognitive rehabilitation group intervention for breast cancer survivors: Results of a randomized clinical trial [published online ahead of print March 10, 2015]. *Psychooncology*. doi: 10.1002/pon.3769
93. Siegel R, Desantis C, Jemal A: Colorectal cancer statistics, 2014. *CA Cancer J Clin* 64:104-117, 2014
94. Simard S, Thewes B, Humphris G, et al: Fear of cancer recurrence in adult cancer survivors: A systematic review of quantitative studies. *J Cancer Surviv* 7:300-322, 2013
95. Boyes AW, Girgis A, Zucca AC, Lecathelinais C: Anxiety and depression among long-term survivors of cancer in Australia: Results of a population-based survey [comment]. *Med J Aust* 191:295, 2009
96. Hoffman KE, McCarthy EP, Recklitis CJ, et al: Psychological distress in long-term survivors of adult-onset cancer: Results from a national survey. *Arch Intern Med* 169:1274-1281, 2009
97. McGinty HL, Goldenberg JL, Jacobsen PB: Relationship of threat appraisal with coping appraisal to fear of cancer recurrence in breast cancer survivors. *Psychooncology* 21:203-210, 2012
98. Mitchell AJ, Ferguson DW, Gill J, et al: Depression and anxiety in long-term cancer survivors compared with spouses and healthy controls: A systematic review and meta-analysis. *Lancet Oncol* 14:721-732, 2013
99. Zainal NZ, Nik-Jaafar NR, Baharudin A, et al: Prevalence of depression in breast cancer survivors: A systematic review of observational studies. *Asian Pac J Cancer Prev* 14:2649-2656, 2013
100. Patel D, Sharpe L, Thewes B, et al: Using the Distress Thermometer and Hospital Anxiety and Depression Scale to screen for psychosocial morbidity in patients diagnosed with colorectal cancer. *J Affect Disord* 131:412-416, 2011
101. Kroenke K, Spitzer RL, Williams JB: The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med* 16:606-613, 2001
102. Spitzer RL, Kroenke K, Williams JB, et al: A brief measure for assessing generalized anxiety disorder: The GAD-7. *Arch Intern Med* 166: 1092-1097, 2006
103. Ewertz M, Jensen AB: Late effects of breast cancer treatment and potentials for rehabilitation. *Acta Oncol* 50:187-193, 2011
104. Ha EH, Cho YK: The mediating effects of self-esteem and optimism on the relationship between quality of life and depressive symptoms of breast cancer patients. *Psychiatry Investig* 11:437-445, 2014
105. Rosen RC, Lane RM, Menza M: Effects of SSRIs on sexual function: A critical review. *J Clin Psychopharmacol* 19:67-85, 1999
106. Speer JJ, Hillenber B, Sugrue DP, et al: Study of sexual functioning determinants in breast cancer survivors. *Breast J* 11:440-447, 2005
107. Casellas-Grau A, Font A, Vives J: Positive psychology interventions in breast cancer. A systematic review. *Psychooncology* 23:9-19, 2014
108. Bower JE: Behavioral symptoms in patients with breast cancer and survivors. *J Clin Oncol* 26: 768-777, 2008
109. Jacobsen PB, Hann DM, Azzarello LM, et al: Fatigue in women receiving adjuvant chemotherapy for breast cancer: Characteristics, course, and correlates. *J Pain Symptom Manage* 18:233-242, 1999
110. Gaston-Johansson F, Fall-Dickson JM, Bakos AB, et al: Fatigue, pain, and depression in pre-autotransplant breast cancer patients. *Cancer Pract* 7:240-247, 1999
111. Meneses-Echavez JF, Gonzalez-Jimenez E, Ramirez-Velez R: Effects of supervised exercise on cancer-related fatigue in breast cancer survivors: A systematic review and meta-analysis [serial online]. *BMC Cancer* 15:77, 2015
112. Gielissen MF, Verhagen CA, Bleijenberg G: Cognitive behaviour therapy for fatigued cancer survivors: Long-term follow-up. *Br J Cancer* 97: 612-618, 2007
113. Duijts SF, Faber MM, Oldenburg HS, et al: Effectiveness of behavioral techniques and physical exercise on psychosocial functioning and health-

related quality of life in breast cancer patients and survivors—a meta-analysis. *Psychooncology* 20:115-126, 2011

114. Kanis JA, McCloskey EV, Powles T, et al: A high incidence of vertebral fracture in women with breast cancer. *Br J Cancer* 79:1179-1181, 1999

115. Chen Z, Maricic M, Bassford TL, et al: Fracture risk among breast cancer survivors: Results from the Women's Health Initiative Observational Study. *Arch Intern Med* 165:552-558, 2005

116. Chen Z, Maricic M, Pettinger M, et al: Osteoporosis and rate of bone loss among postmenopausal survivors of breast cancer. *Cancer* 104:1520-1530, 2005

117. Lindsey AM, Gross G, Twiss J, et al: Postmenopausal survivors of breast cancer at risk for osteoporosis: Nutritional intake and body size. *Cancer Nurs* 25:50-56, 2002

118. Gralow JR, Biermann JS, Farooki A, et al: NCCN Task Force Report: Bone Health in Cancer Care. *J Natl Compr Canc Netw* 11:S1-S50; quiz S51, 2013 (suppl 3)

119. Johnell O, Kanis JA, Oden A, et al: Predictive value of BMD for hip and other fractures. *J Bone Miner Res* 20:1185-1194, 2005

120. Kanis JA, Borgstrom F, De Laet C, et al: Assessment of fracture risk. *Osteoporos Int* 16:581-589, 2005

121. Feskanich D, Willett W, Colditz G: Walking and leisure-time activity and risk of hip fracture in postmenopausal women. *JAMA* 288:2300-2306, 2002

122. Moyer VA: Prevention of falls in community-dwelling older adults: US Preventive Services Task Force recommendation statement. *Ann Intern Med* 157:197-204, 2012

123. Cosman F, de Beur SJ, LeBoff MS, et al: Clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int* 25:2359-2381, 2014

124. Chapuy MC, Arlot ME, Duboeuf F, et al: Vitamin D3 and calcium to prevent hip fractures in the elderly women. *N Engl J Med* 327:1637-1642, 1992

125. Dawson-Hughes B, Harris SS, Krall EA, et al: Effect of calcium and vitamin D supplementation on bone density in men and women 65 years of age or older. *N Engl J Med* 337:670-676, 1997

126. Murad MH, Elamin KB, Abu Elnour NO, et al: Clinical review: The effect of vitamin D on falls: A systematic review and meta-analysis. *J Clin Endocrinol Metab* 96:2997-3006, 2011

127. Greenspan SL, Nelson JB, Trump DL, et al: Effect of once-weekly oral alendronate on bone loss in men receiving androgen deprivation therapy for prostate cancer: A randomized trial. *Ann Intern Med* 146:416-424, 2007

128. Van Poznak C: Managing bone mineral density with oral bisphosphonate therapy in women with breast cancer receiving adjuvant aromatase inhibition [serial online]. *Breast Cancer Res* 12:110, 2010

129. Van Poznak C, Hannon RA, Mackey JR, et al: Prevention of aromatase inhibitor-induced bone loss using risenedronate: The SABRE trial. *J Clin Oncol* 28:967-975, 2010

130. Baum M, Budzar AU, Cuzick J, et al: Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early breast cancer: First results of the ATAC randomised trial. *Lancet* 359:2131-2139, 2002

131. Stubblefield MD, Keole N: Upper body pain and functional disorders in patients with breast cancer. *PM R* 6:170-183, 2014

132. Harrington S, Padua D, Battaglini C, et al: A survey of joint and muscle aches, pain, and stiffness comparing women with and without breast cancer. *J Pain Symptom Manage* 46:523-535, 2013

133. Fenlon D, Addington-Hall JM, O'Callaghan AC, et al: A survey of joint and muscle aches, pain, and stiffness comparing women with and without breast cancer. *J Pain Symptom Manage* :523-5, 2013

134. Karki A, Simonen R, Malkia E, et al: Impairments, activity limitations and participation restrictions 6 and 12 months after breast cancer operation. *J Rehabil Med* 37:180-188, 2005

135. Kakuda JT, Stuntz M, Trivedi V, et al: Objective assessment of axillary morbidity in breast cancer treatment. *Am Surg* 65:995-998, 1999

136. Kuehn T, Klaus W, Darsow M, et al: Long-term morbidity following axillary dissection in breast cancer patients—clinical assessment, significance for life quality and the impact of demographic, oncologic and therapeutic factors. *Breast Cancer Res Treat* 64:275-286, 2000

137. Crew KD, Greenlee H, Capodice J, et al: Prevalence of joint symptoms in postmenopausal women taking aromatase inhibitors for early-stage breast cancer. *J Clin Oncol* 25:3877-3883, 2007

138. Henry NL, Azzouz F, Desta Z, et al: Predictors of aromatase inhibitor discontinuation as a result of treatment-emergent symptoms in early-stage breast cancer. *J Clin Oncol* 30:936-942, 2012

139. Hershman DL, Kushi LH, Shao T, et al: Early discontinuation and nonadherence to adjuvant hormonal therapy in a cohort of 8,769 early-stage breast cancer patients. *J Clin Oncol* 28:4120-4128, 2010

140. Barron TI, Cahir C, Sharp L, et al: A nested case-control study of adjuvant hormonal therapy persistence and compliance, and early breast cancer recurrence in women with stage I-III breast cancer. *Br J Cancer* 109:1513-1521, 2013

141. De Groef A, Van Kampen M, Dieltjens E, et al: Effectiveness of postoperative physical therapy for upper-limb impairments after breast cancer treatment: A systematic review. *Arch Phys Med Rehabil* 96:1140-1153, 2015

142. Loh SY, Musa AN: Methods to improve rehabilitation of patients following breast cancer surgery: A review of systematic reviews. *Breast Cancer (Dove Med Press)* 7:81-98, 2015

143. Irwin ML, Cartmel B, Gross CP, et al: Randomized exercise trial of aromatase inhibitor-induced arthralgia in breast cancer survivors. *J Clin Oncol* 33:1104-1111, 2015

144. Crew KD, Capodice JL, Greenlee H, et al: Randomized, blinded, sham-controlled trial of acupuncture for the management of aromatase inhibitor-associated joint symptoms in women with early-stage breast cancer. *J Clin Oncol* 28:1154-1160, 2010

145. Andersen KG, Kehlet H: Persistent pain after breast cancer treatment: A critical review of risk factors and strategies for prevention. *J Pain* 12:725-746, 2011

146. Gartner R, Jensen MB, Nielsen J, et al: Prevalence of and factors associated with persistent pain following breast cancer surgery. *JAMA* 302:1985-1992, 2009

147. Pachman DR, Barton DL, Watson JC, et al: Chemotherapy-induced peripheral neuropathy: Prevention and treatment. *Clin Pharmacol Ther* 90:377-387, 2011

148. Steegers MA, Snik DM, Verhagen AF, et al: Only half of the chronic pain after thoracic surgery shows a neuropathic component. *J Pain* 9:955-961, 2008

149. Garcia MK, McQuade J, Haddad R, et al: Systematic review of acupuncture in cancer care: A synthesis of the evidence. *J Clin Oncol* 31:952-960, 2013

150. Mishra SI, Scherer RW, Geigle PM, et al: Exercise interventions on health-related quality of life for cancer survivors [serial online]. *Cochrane Database Syst Rev* (8):CD007566, 2012

151. Courneya KS, McKenzie DC, Mackey JR, et al: Subgroup effects in a randomized trial of different types and doses of exercise during breast cancer chemotherapy. *Br J Cancer* 111:1718-1725, 2014

152. Smith E, Pang H, Cirrincione C, et al: Effect of duloxetine on pain, function, and quality of life among patients with chemotherapy-induced painful peripheral neuropathy: A randomized clinical trial. *JAMA* 309:1359-1367, 2013

153. Kort JD, Eisenberg ML, Millheiser LS, et al: Fertility issues in cancer survivorship. *CA Cancer J Clin* 64:118-134, 2014

154. Ganz PA, Land SR, Geyer CE Jr., et al: Menstrual history and quality-of-life outcomes in women with node-positive breast cancer treated with adjuvant therapy on the NSABP B-30 trial. *J Clin Oncol* 29:1110-1116, 2011

155. Rosen MP, Sternfeld B, Schuh-Huerta SM, et al: Antral follicle count: Absence of significant midlife decline. *Fertil Steril* 94:2182-2185, 2010

156. Sadovsky R, Basson R, Krychman M, et al: Cancer and sexual problems. *J Sex Med* 7:349-373, 2010

157. Berkey FJ: Managing the adverse effects of radiation therapy. *Am Fam Physician* 82:381-388, 394, 2010

158. Mazzeo S, Hutton B, Ibrahim MF, et al: Management of urogenital atrophy in breast cancer patients: A systematic review of available evidence from randomized trials. *Breast Cancer Res Treat* 152:1-8, 2015

159. Pruthi S, Simon JA, Early AP: Current overview of the management of urogenital atrophy in women with breast cancer. *Breast J* 17:403-408, 2011

160. Kendall A, Dowsett M, Folkler E, et al: Caution: Vaginal estradiol appears to be contraindicated in postmenopausal women on adjuvant aromatase inhibitors. *Ann Oncol* 17:584-587, 2006

161. Rowland JH, Meyerowitz BE, Crespi CM, et al: Addressing intimacy and partner communication after breast cancer: A randomized controlled group intervention. *Breast Cancer Res Treat* 118:99-111, 2009

162. Murthy V, Chamberlain RS: Menopausal symptoms in young survivors of breast cancer: A growing problem without an ideal solution. *Cancer Control* 19:317-329, 2012

163. Kaplan M, Mahon S, Cope D, et al: Putting evidence into practice: Evidence-based interventions for hot flashes resulting from cancer therapies. *Clin J Oncol Nurs* 15:149-157, 2011

164. Jin Y, Desta Z, Stearns V, et al: CYP2D6 genotype, antidepressant use, and tamoxifen metabolism during adjuvant breast cancer treatment. *J Natl Cancer Inst* 97:30-39, 2005

165. Burstein HJ, Prestrud AA, Seidenfeld J, et al: American Society of Clinical Oncology clinical practice guideline: Update on adjuvant endocrine therapy for women with hormone receptor-positive breast cancer. *J Clin Oncol* 28:3784-3796, 2010

166. Loprinzi CL, Sloan J, Stearns V, et al: Newer antidepressants and gabapentin for hot flashes: An individual patient pooled analysis. *J Clin Oncol* 27:2831-2837, 2009
167. Bordeleau L, Pritchard KI, Loprinzi CL, et al: Multicenter, randomized, cross-over clinical trial of venlafaxine versus gabapentin for the management of hot flashes in breast cancer survivors. *J Clin Oncol* 28:5147-5152, 2010
168. Pandya KJ, Morrow GR, Roscoe JA, et al: Gabapentin for hot flashes in 420 women with breast cancer: A randomised double-blind placebo-controlled trial. *Lancet* 366:818-824, 2005
169. Clayden JR, Bell JW, Pollard P: Menopausal flushing: Double-blind trial of a nonhormonal medication. *BMJ* 1:409-412, 1974
170. Chiu HY, Shyu YK, Chang PC, Tsai PS: Effects of acupuncture on menopause-related symptoms in breast cancer survivors: A meta-analysis of randomized controlled trials [published online ahead of print June 3, 2015]. *Cancer Nurs*. doi: [10.1097/NCC.0000000000000278](https://doi.org/10.1097/NCC.0000000000000278)
171. Bordeleau L, Pritchard K, Goodwin P, et al: Therapeutic options for the management of hot flashes in breast cancer survivors: An evidence-based review. *Clin Ther* 29:230-241, 2007
172. Chan DS, Vieira AR, Aune D, et al: Body mass index and survival in women with breast cancer—systematic literature review and meta-analysis of 82 follow-up studies. *Ann Oncol* 25:1901-1914, 2014
173. Protani M, Coory M, Martin JH: Effect of obesity on survival of women with breast cancer: Systematic review and meta-analysis. *Breast Cancer Res Treat* 123:627-635, 2010
174. McTiernan A, Irwin M, Vongruenigen V: Weight, physical activity, diet, and prognosis in breast and gynecologic cancers. *J Clin Oncol* 28:4074-4080, 2010
175. Wei EK, Wolin KY, Colditz GA: Time course of risk factors in cancer etiology and progression. *J Clin Oncol* 28:4052-4057, 2010
176. Patterson RE, Cadmus LA, Emond JA, et al: Physical activity, diet, adiposity and female breast cancer prognosis: A review of the epidemiologic literature. *Maturitas* 66:5-15, 2010
177. Kellen E, Vansant G, Christiaens MR, et al: Lifestyle changes and breast cancer prognosis: A review. *Breast Cancer Res Treat* 114:13-22, 2009
178. Friedenreich CM: Physical activity and breast cancer: Review of the epidemiologic evidence and biologic mechanisms. *Recent Results Cancer Res* 188:125-139, 2011
179. Barbarks E, Mooic M, Broore L, Cheifetz O: Effects of physical activity on cancer survival: A systematic review. *Physiother Can* 62:25-34, 2010
180. Carmichael AR, Daley AJ, Rea DW, et al: Physical activity and breast cancer outcome: A brief review of evidence, current practice and future direction. *Eur J Surg Oncol* 36:1139-1148, 2010
181. Speed-Andrews AE, Courneya KS: Effects of exercise on quality of life and prognosis in cancer survivors. *Curr Sports Med Rep* 8:176-181, 2009
182. Irwin ML, Mayne ST: Impact of nutrition and exercise on cancer survival. *Cancer J* 14:435-441, 2008
183. Lahart IM, Metsios GS, Nevill AM, et al: Physical activity, risk of death and recurrence in breast cancer survivors: A systematic review and meta-analysis of epidemiological studies. *Acta Oncol* 54:635-654, 2015
184. Ballard-Barbash R, Friedenreich CM, Courneya KS, et al: Physical activity, biomarkers, and disease outcomes in cancer survivors: A systematic review. *J Natl Cancer Inst* 104:815-840, 2012
185. Binkley JM, Harris SR, Levangie PK, et al: Patient perspectives on breast cancer treatment side effects and the prospective surveillance model for physical rehabilitation for women with breast cancer. *Cancer* 118:2207-2216, 2012
186. Cappiello M, Cunningham RS, Knobf MT, et al: Breast cancer survivors: Information and support after treatment. *Clin Nurs Res* 16:278-293, 2007; discussion 294-301
187. Bloom JR, Stewart SL, Chang S, et al: Then and now: Quality of life of young breast cancer survivors. *Psychooncology* 13:147-160, 2004
188. Blank TO, Bellizzi KM: A gerontologic perspective on cancer and aging. *Cancer* 112:2569-2576, 2008
189. Howard-Anderson J, Ganz PA, Bower JE, et al: Quality of life, fertility concerns, and behavioral health outcomes in younger breast cancer survivors: A systematic review. *J Natl Cancer Inst* 104:386-405, 2012
190. Stricker CT, Jacobs LA, Risendal B, et al: Survivorship care planning after the Institute of Medicine recommendations: How are we faring? *J Cancer Surviv* 5:358-370, 2011
191. Brennan ME, Gormally JF, Butow P, et al: Survivorship care plans in cancer: A systematic review of care plan outcomes. *Br J Cancer* 111:1899-1908, 2014
192. Mayer DK, Birken SA, Check DK, et al: Summing it up: An integrative review of studies of cancer survivorship care plans (2006-2013). *Cancer* 121:978-996, 2015
193. Grunfeld E, Julian JA, Pond G, et al: Evaluating survivorship care plans: Results of a randomized, clinical trial of patients with breast cancer. *J Clin Oncol* 29:4755-4762, 2011
194. Brothers BM, Easley A, Salani R, et al: Do survivorship care plans impact patients' evaluations of care? A randomized evaluation with gynecologic oncology patients. *Gynecol Oncol* 129:554-558, 2013
195. Hershman DL, Greenlee H, Awad D, et al: Randomized controlled trial of a clinic-based survivorship intervention following adjuvant therapy in breast cancer survivors. *Breast Cancer Res Treat* 138:795-806, 2013
196. Ezendam NP, Nicolaije KA, Kruitwagen RF, et al: Survivorship Care Plans to inform the primary care physician: Results from the ROGY care pragmatic cluster randomized controlled trial. *J Cancer Surviv* 8:595-602, 2014
197. American Cancer Society. The American Cancer Society web page. cancer.org. Accessed May 15, 2015
198. Irwin ML, McTiernan A, Bernstein L, et al: Physical activity levels among breast cancer survivors. *Med Sci Sports Exerc* 36:1484-1491, 2004
199. Demark-Wahnefried W, Rogers LO, Alfano CM, et al: Practical clinical interventions for diet, physical activity, and weight control in cancer survivors. *CA Cancer J Clin* 65:167-189, 2015
200. Ligibel JA, Alfano CM, Courneya KS, et al: American Society of Clinical Oncology position statement on obesity and cancer. *J Clin Oncol* 32:3568-3574, 2014
201. US Preventive Services Task Force. Obesity in Adults: Screening and Management, June 2012. [uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/obesity-in-adults-screening-and-management#references](https://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/obesity-in-adults-screening-and-management#references). Accessed August 17, 2015
202. Underwood JM, Townsend JS, Stewart SL, et al: Surveillance of demographic characteristics and health behaviors among adult cancer survivors—Behavioral Risk Factor Surveillance System, United States, 2009. *MMWR Surveill Summ* 61:1-23, 2012
203. Schmid D, Leitzmann MF: Association between physical activity and mortality among breast cancer and colorectal cancer survivors: A systematic review and meta-analysis. *Ann Oncol* 25:1293-1311, 2014
204. Zhao G, Li C, Okoro CA, et al: Trends in modifiable lifestyle-related risk factors following diagnosis in breast cancer survivors. *J Cancer Surviv* 7:563-569, 2013
205. Blanchard CM, Courneya KS, Stein K: Cancer survivors' adherence to lifestyle behavior recommendations and associations with health-related quality of life: Results from the American Cancer Society's SCS-II. *J Clin Oncol* 26:2198-2204, 2008
206. Forbes CC, Blanchard CM, Mummery WK, et al: Prevalence and correlates of strength exercise among breast, prostate, and colorectal cancer survivors. *Oncol Nurs Forum* 42:118-127, 2015
207. Irwin ML, McTiernan A, Manson JE, et al: Physical activity and survival in postmenopausal women with breast cancer: Results from the Women's Health Initiative. *Cancer Prev Res (Phila)* 4:522-529, 2011
208. Kroenke CH, Fung TT, Hu FB, et al: Dietary patterns and survival after breast cancer diagnosis. *J Clin Oncol* 2005;23:9295-9303. Kwan ML, Weltzien E, Kushi LH, Castillo A, Slattery ML, Caan BJ: Dietary patterns and breast cancer recurrence and survival among women with early-stage breast cancer. *J Clin Oncol* 27:919-926, 2009
209. Vrieling A, Buck K, Seibold P, et al: Dietary patterns and survival in German postmenopausal breast cancer survivors. *Br J Cancer* 108:188-192, 2013
210. Chlebowski RT, Blackburn GL, Thomson CA, et al: Dietary fat reduction and breast cancer outcome: Interim efficacy results from the Women's Intervention Nutrition Study. *J Natl Cancer Inst* 98:1767-1776, 2006
211. Pierce JP, Natarajan L, Caan BJ, et al: Influence of a diet very high in vegetables, fruit, and fiber and low in fat on prognosis following treatment for breast cancer: The Women's Healthy Eating and Living (WHEL) randomized trial. *JAMA* 298:289-298, 2007
212. American Cancer Society. Alcohol Use and Cancer. cancer.org/cancer/cancercauses/dietandphysicalactivity/alcohol-use-and-cancer. Accessed January 6, 2015
213. McGuire S. US Department of Agriculture and US Department of Health and Human Services, Dietary Guidelines for Americans, 2010. 7th edition, Washington, DC: US Government Printing Office, January 2011. *Adv Nutr* 2011;2:293-294
214. Kwan ML, Kushi LH, Weltzien E, et al: Alcohol consumption and breast cancer recurrence and survival among women with early-stage breast cancer: The life after cancer epidemiology study. *J Clin Oncol* 28:4410-4416, 2010
215. Baan R, Straif K, Grosse Y, et al: Carcinogenicity of alcoholic beverages. *Lancet Oncol* 8:292-293, 2007
216. Nelson DE, Jarman DW, Rehm J, et al: Alcohol-attributable cancer deaths and years of potential life lost in the United States. *Am J Public Health* 103:641-648, 2013
217. Scoccianti C, Lauby-Secretan B, Bello PY, et al: Female breast cancer and alcohol consumption: A

review of the literature. *Am J Prev Med* 46(3 suppl 1): S16-S25, 2014

218. Terry MB, Zhang FF, Kabat G, et al: Lifetime alcohol intake and breast cancer risk. *Ann Epidemiol* 16:230-240, 2006

219. Berube S, Lemieux J, Moore L, et al: Smoking at time of diagnosis and breast cancer-specific survival: New findings and systematic review with meta-analysis [serial online]. *Breast Cancer Res* 16:R42, 2014

220. Nayan S, Gupta MK, Strychowski JE, et al: Smoking cessation interventions and cessation rates in the oncology population: An updated systematic review and meta-analysis. *Otolaryngol Head Neck Surg* 149:200-211, 2013

221. Boekhout AH, Maunsell E, Pond GR, et al: A survivorship care plan for breast cancer survivors: Extended results of a randomized clinical trial [published online ahead of print April 21, 2015]. *J Cancer Surviv* doi: [10.1007/s11764-015-0443-1](https://doi.org/10.1007/s11764-015-0443-1)

222. Mao JJ, Bowman MA, Stricker CT, et al: Delivery of survivorship care by primary care physicians: The perspective of breast cancer patients. *J Clin Oncol* 27:933-938, 2009

223. Shulman LN, Jacobs LA, Greenfield S, et al: Cancer care and cancer survivorship care in the United States: Will we be able to care for these patients in the future? *J Oncol Pract* 5:119-123, 2009

224. Travis LB, Yahalom J: Cancer Survivorship. Preface. *Hematol Oncol Clin North Am* 22:xi-xii, 2008

225. Children's Oncology Group: Long-Term Follow-Up Guidelines for Survivors of Childhood, Adolescent, and Young Adult Cancers. survivorship-guidelines.org. Accessed May 15, 2015

226. Grunfeld E, Levine MN, Julian JA, et al: Randomized trial of long-term follow-up for early-stage breast cancer: A comparison of family physician versus specialist care. *J Clin Oncol* 24:848-855, 2006

227. Grunfeld E, Fitzpatrick R, Mant D, et al: Comparison of breast cancer patient satisfaction with follow-up in primary care versus specialist care: Results from a randomized controlled trial. *Br J Gen Pract* 49:705-710, 1999

228. Dawes AJ, Hemmelgarn M, Nguyen DK, et al: Are primary care providers prepared to care for survivors of breast cancer in the safety net? *Cancer* 121: 1249-1256, 2015

229. Palmer SC, Stricker CT, Panzer SL, et al: Outcomes and satisfaction after delivery of a breast cancer survivorship care plan: Results of a multicenter trial [serial online]. *J Oncol Pract* 11: e222-e229, 2015

230. Kim Y, Kashy DA, Spillers RL, et al: Needs assessment of family caregivers of cancer survivors: Three cohorts comparison. *Psychooncology* 19: 573-582, 2010

231. Kim Y, Spillers RL, Hall DL: Quality of life of family caregivers 5 years after a relative's cancer diagnosis: Follow-up of the national quality of life

survey for caregivers. *Psychooncology* 21:273-281, 2012

232. Given BA, Given CW, Sherwood PR: Family and caregiver needs over the course of the cancer trajectory. *J Support Oncol* 10:57-64, 2012

233. National Center for Chronic Disease Prevention and Health Promotion. Centers for Disease Control and Prevention, United States Cancer Statistics: 1999-2011 Incidence and Mortality [web-based report]. nccd.cdc.gov/uscs/. Accessed April 1, 2015

234. American Cancer Society. *Cancer Facts & Figures for African Americans 2013-2014*. Atlanta, GA: American Cancer Society; 2013

235. National Cancer Institute; Surveillance, Epidemiology, and End Results (SEER) Program. *SEER Cancer Statistics Review, 1975-2011*. seer.cancer.gov/archive/csr/1975_2011/. Accessed April 1, 2015

236. Mead H, Cartwright-Smith L, Jones K, Ramos C, Woods K, Siegel B. *Racial and Ethnic Disparities in US Health Care: A Chartbook*. New York, NY: The Commonwealth Fund; 2008

237. American Geriatrics Society Expert Panel on the Care of Older Adults With Multimorbidity. Patient-centered care for older adults with multiple chronic conditions: A stepwise approach from the American Geriatrics Society. *J Am Geriatr Soc* 60:1957-1968, 2012

Affiliations

Carolyn D. Runowicz, Herbert Wertheim College of Medicine, Florida International University; Karen S. Henry, Sylvester Cancer Center at the University of Miami, Miami, FL; Corinne R. Leach, Rebecca L. Cowens-Alvarado, Rachel S. Cannady, and Samuel J. LaMonte, American Cancer Society, Atlanta, GA; N. Lynn Henry, University of Michigan, Comprehensive Cancer Center, Ann Arbor, MI; Heather T. Mackey, Oncology Nursing Society, Pittsburgh; Linda A. Jacobs, Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA; Mandi L. Pratt-Chapman, The George Washington University Cancer Institute, Washington, DC; Stephen B. Edge, Baptist Cancer Center, Memphis, TN; Arti Hurria, City of Hope, Duarte; Patricia A. Ganz, Schools of Medicine and Public Health, University of California, Los Angeles, CA; Lawrence B. Marks, University of North Carolina, Chapel Hill, NC; Ellen Warner, University of Toronto, Sunnybrook Odette Cancer Centre, Toronto, Ontario, Canada; and Gary H. Lyman, Hutchinson Institute for Cancer Outcomes Research, Fred Hutchinson Cancer Research Center, Seattle, WA doi: [10.3322/caac.21319](https://doi.org/10.3322/caac.21319). Available online at cancerjournal.com.

Support

This journal article was supported, in part, by Cooperative Agreement 5U55DP003054 from the Centers for Disease Control and Prevention. No industry funding was used to support this work.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or jco.ascopubs.org/site/ifc.

Carolyn D. Runowicz

Honoraria: Hologic, UpToDate, Audio Digest, Pfizer

Consulting or Advisory Role: Pfizer

Patents, Royalties, Other Intellectual Property: Patent for "High efficacy anticancer drug delivery, release and nanoscale heat treatment of cancerous cells" (Inst)

Travel, Accommodations, Expenses: Hologic, Pfizer

Corinne R. Leach

No relationship to disclose

N.Lynn Henry

Research Funding: Sanofi, BioMarin, Celldex

Travel, Accommodations, Expenses: Celldex

Karen S. Henry

No relationship to disclose

Heather T. Mackey

Consulting or Advisory Role: Oncology Nursing Society

Rebecca L. Cowens-Alvarado

No relationship to disclose

Rachel S. Cannady

No relationship to disclose

Mandi L. Pratt-Chapman

Consulting or Advisory Role: Pfizer

Research Funding: Genentech (Inst), Amgen (Inst), Takeda (Inst)

Patents, Royalties, Other Intellectual Property: The GW Cancer Institute retains intellectual property for products created by the GW Cancer Institute and the Center for Advancing Health (Prepared Patient TM, Aftershock TM). (Inst)

Stephen B. Edge

No relationship to disclose

Linda A. Jacobs

No relationship to disclose

Arti Hurria

Consulting or Advisory Role: GTx, Seattle Genetics, Boehringer Ingelheim

Research Funding: GlaxoSmithKline (Inst), Celgene (Inst)

Lawrence B. Marks

Research Funding: Elekta, Morphormics, Accuray, Sun Nuclear, United Health International (Inst)

Travel, Accommodations, Expenses: Varian, Elekta

Samuel J. LaMonte

No relationship to disclose

Ellen Warner

No relationship to disclose

Gary H. Lyman

Research Funding: Amgen (Inst)

Patricia A. Ganz

Leadership: Intrinsic LifeSciences (I)

Stock or Other Ownership: xenon (I), Intrinsic LifeSciences (I), Silarus Therapeutics (I), Merganser Biotech (I), Teva, Novartis, Merck, Johnson & Johnson, Pfizer, GlaxoSmithKline, Abbott Laboratories

Consulting or Advisory Role: Keryx (I), Merganser Biotech (I), Silarus Therapeutics (I), InformedDNA

Research Funding: Breast Cancer Research Foundation, Susan G. Komen for the Cure, Keryx (I)

Patents, Royalties, Other Intellectual Property: Related to iron metabolism and the anemia of chronic disease (I), Up-to-Date royalties for section editor on survivorship

Travel, Accommodations, Expenses: Intrinsic LifeSciences (I), Keryx (I)

Acknowledgment

We thank Dr Catherine Alfano, Dr Neelima Denduluri, Dr Lewis Foxhall, Dr Ted Gansler, Dr Michael Jefford, Dr Mariana Chavez Mac Gregor, Thomas Oliver, Christina Lacchetti, Dr Marcus Plescia, Dr Robert Smith, Alyssa Troeschel, Dr Richard Wender, and the Clinical Practice Guidelines Committee for their contributions, thoughtful reviews, and insightful comments on this guideline document. In addition, we thank the following members of the Breast Cancer Survivorship Care Expert Workgroup for their contribution to the development of the Breast Cancer Survivorship Care Guideline: Dr Carmel Cohen, Dr Stephen B. Edge, Dr Patricia A. Ganz, Dr Karen S. Henry, Dr N. Lynn Henry, Dr Arti Hurria, Dr Linda A. Jacobs, Dr Samuel J. LaMonte, Susan Leigh, Dr Heather T. Mackey, Dr Lawrence B. Marks, Dr Carolyn D. Runowicz, Dr Debbie Saslow, Dr Eddie Turner, and Dr Ellen Warner.